



π^0 - η - η' mixing from $V \rightarrow P\gamma$ and $P \rightarrow V\gamma$ decays

Rafel Escribano
UAB and IFAE-BIST

QCHS 2021

The XIVth Quark confinement
and the Hadron spectrum conference

August, 2021
Stavanger (Norway)



This talk is dedicated to Simon Eidelman

What's the goal of this analysis?

To **estimate** the **admixtures** of the η and η' mesons to the physical π^0

$$|\pi^0\rangle = |\pi_3\rangle + \epsilon |\eta\rangle + \epsilon' |\eta'\rangle$$

where π_3 is the **$I_3=0$ state** of the pseudoscalar **isospin triplet**

The responsible of this **mixing** is **isospin breaking**

T. Feldmann, P. Kroll and B. Stech, Phys. Lett. B449 (1999) 339

$$\epsilon = 1.4 \% \qquad \epsilon' = 0.37 \%$$

η - η' mixing: a reminder

octet-singlet basis

$$|\eta\rangle = \cos \theta_P |\eta_8\rangle - \sin \theta_P |\eta_0\rangle \qquad |\eta_8\rangle = \frac{1}{\sqrt{6}} |u\bar{u} + d\bar{d} - 2s\bar{s}\rangle$$

$$|\eta'\rangle = \sin \theta_P |\eta_8\rangle + \cos \theta_P |\eta_0\rangle \qquad |\eta_0\rangle = \frac{1}{\sqrt{3}} |u\bar{u} + d\bar{d} + s\bar{s}\rangle$$

quark-flavour basis

$$|\eta\rangle = \cos \phi_P |\eta_{\text{NS}}\rangle - \sin \phi_P |\eta_{\text{S}}\rangle \qquad |\eta_{\text{NS}}\rangle = \frac{1}{\sqrt{2}} |u\bar{u} + d\bar{d}\rangle$$

$$|\eta'\rangle = \sin \phi_P |\eta_{\text{NS}}\rangle + \cos \phi_P |\eta_{\text{S}}\rangle \qquad |\eta_{\text{S}}\rangle = |s\bar{s}\rangle$$

$$\theta_P = \phi_P - \arctan \sqrt{2} \simeq \phi_P - 54.7^\circ$$

η - η' is heavily influenced by the $\text{U}(1)_\text{A}$ of QCD

Previous estimates

Kroll based on the FKS scheme

$$\begin{aligned}\epsilon(z) &= \cos \phi \left[\frac{1}{2} \frac{m_{dd}^2 - m_{uu}^2}{m_{\eta}^2 - m_{\pi^0}^2} + z \right] \\ \epsilon'(z) &= \sin \phi \left[\frac{1}{2} \frac{m_{dd}^2 - m_{uu}^2}{m_{\eta'}^2 - m_{\pi^0}^2} + z \right]\end{aligned} \quad \longrightarrow \quad \begin{aligned}\phi &= 39.3^\circ \\ \hat{\epsilon} &= \epsilon(z=0) = (1.7 \pm 0.2)\% \\ \hat{\epsilon}' &= \epsilon'(z=0) = (0.4 \pm 0.1)\%\end{aligned}$$

P. Kroll, Mod. Phys. Lett. A20 (2005) 2667

Escribano et al. based on ChPT with resonances

$$\begin{aligned}\epsilon_{\pi\eta} &= c\phi_{\eta\eta'} \frac{m_{K^0}^2 - m_{K^+}^2 - m_{\pi^0}^2 + m_{\pi^+}^2}{m_{\eta}^2 - m_{\pi^-}^2} \left[1 - \frac{m_{\eta}^2 - m_{\pi^-}^2}{M_S^2} \right] \\ \epsilon_{\pi\eta'} &= s\phi_{\eta\eta'} \frac{m_{K^0}^2 - m_{K^+}^2 - m_{\pi^0}^2 + m_{\pi^+}^2}{m_{\eta'}^2 - m_{\pi^-}^2} \left[1 - \frac{m_{\eta'}^2 - m_{\pi^-}^2}{M_S^2} \right]\end{aligned} \quad \longrightarrow \quad \begin{aligned}\phi_{\eta\eta'} &= (41.4 \pm 0.5)^\circ \\ (9.8 \pm 0.3) &\times 10^{-3} \\ (2.5 \pm 1.5) &\times 10^{-4}\end{aligned}$$

R. Escribano, S. González-Solís and P. Roig, Phys. Rev. D94 (2016) 034008

The experimental data

$V \rightarrow P\gamma$ and $P \rightarrow V\gamma$ decays is

the most **extensive** and **exhaustive** set of data

| | Transition | Γ_{exp} (keV) | |
|--------------------|----------------------------------|-----------------------------|------------|
| | $\rho^0 \rightarrow \eta\gamma$ | 44 ± 3 | 7% |
| | $\rho^0 \rightarrow \pi^0\gamma$ | 69 ± 9 | 13% |
| | $\rho^+ \rightarrow \pi^+\gamma$ | 67 ± 7 | 10% |
| | $\omega \rightarrow \eta\gamma$ | 3.8 ± 0.3 | 8% |
| SND 2013 | $\omega \rightarrow \pi^0\gamma$ | 713 ± 20 | 3% |
| SND 2000 | $\phi \rightarrow \eta\gamma$ | 55.4 ± 1.1 | 2% |
| KLOE 2007 | $\phi \rightarrow \eta'\gamma$ | 0.26 ± 0.01 | 4% |
| SND 2000 | $\phi \rightarrow \pi^0\gamma$ | 5.5 ± 0.2 | 4% |
| BESIII 2018 | $\eta' \rightarrow \rho^0\gamma$ | 57 ± 3 | 5% |
| BES 2019 | $\eta' \rightarrow \omega\gamma$ | 5.1 ± 0.3 | 6% |
| | $K^{*0} \rightarrow K^0\gamma$ | 116 ± 10 | 9% |
| | $K^{*+} \rightarrow K^+\gamma$ | 46 ± 4 | 9% |

The theoretical model

The **most general** $SU(3)_F$ -**symmetric** effective Lagrangian consistent with **Lorentz**, **P** and **C** invariance

$$\mathcal{L}_{VP\gamma} = g_e \epsilon_{\mu\nu\alpha\beta} \partial^\mu A^\nu \text{Tr}[Q (\partial^\alpha V^\beta P + P \partial^\alpha V^\beta)]$$

supplemented with **conventional quark model** ideas to introduce **flavour** and **isospin breaking**

- **Magnetic dipole transitions**

$$\mu_q = e_q / 2m_q \quad 1 - s_e \equiv \bar{m}/m_s$$

- **Relative overlap** between the **P** and **V** wavefunctions

OZI-rule: $Z_\pi = \langle \pi | \omega_{NS} \rangle = \langle \pi | \rho \rangle \quad Z_{NS} = \langle \eta_{NS} | \omega_{NS} \rangle = \langle \eta_{NS} | \rho \rangle \quad Z_S = \langle \eta_S | \omega_S \rangle$

The theoretical model

π^0 - η - η' mixing in the quark-flavour basis

$$\begin{pmatrix} \pi^0 \\ \eta \\ \eta' \end{pmatrix} = \begin{pmatrix} 1 & \epsilon_{12} & \epsilon_{13} \\ -\epsilon_{12} \cos \phi_{23} + \epsilon_{13} \sin \phi_{23} & \cos \phi_{23} & -\sin \phi_{23} \\ -\epsilon_{13} \cos \phi_{23} - \epsilon_{12} \sin \phi_{23} & \sin \phi_{23} & \cos \phi_{23} \end{pmatrix} \begin{pmatrix} \pi_3 \\ \eta_{\text{NS}} \\ \eta_{\text{S}} \end{pmatrix}$$

To compare to Kroll's results:

$$\begin{pmatrix} \epsilon_{12} \\ \epsilon_{13} \end{pmatrix} = \begin{pmatrix} \cos \phi_P & \sin \phi_P \\ -\sin \phi_P & \cos \phi_P \end{pmatrix} \begin{pmatrix} \epsilon \\ \epsilon' \end{pmatrix}$$

To compare to Escribano et al.'s results:

$$\begin{pmatrix} \epsilon_{12} \\ \epsilon_{13} \end{pmatrix} = \frac{1}{\sqrt{3}} \begin{pmatrix} \cos \theta_P - \sqrt{2} \sin \theta_P & \sin \theta_P + \sqrt{2} \cos \theta_P \\ -\sin \theta_P - \sqrt{2} \cos \theta_P & \cos \theta_P - \sqrt{2} \sin \theta_P \end{pmatrix} \begin{pmatrix} \epsilon_{\pi\eta} \\ \epsilon_{\pi\eta'} \end{pmatrix}$$

The theoretical model

Couplings of the enhanced phenomenological model

$$\begin{aligned}g_{\rho^0\pi^0\gamma} &= g\left(\frac{1}{3} + \epsilon_{12}Z_{\text{NS}}\right), \quad g_{\rho^+\pi^+\gamma} = g\frac{z_+}{3}, \\g_{\rho^0\eta\gamma} &= g\left[\left(Z_{\text{NS}} - \frac{\epsilon_{12}}{3}\right)c\phi_{23} + \frac{\epsilon_{13}}{3}s\phi_{23}\right], \\g_{\omega\pi^0\gamma} &= g\left[\left(1 + \frac{\epsilon_{12}}{3}Z_{\text{NS}}\right)c\phi_V + \frac{2}{3}z_S\frac{\bar{m}}{m_s}\epsilon_{13}s\phi_V\right], \\g_{\eta'\rho^0\gamma} &= g\left[\left(Z_{\text{NS}} - \frac{\epsilon_{12}}{3}\right)s\phi_{23} - \frac{\epsilon_{13}}{3}c\phi_{23}\right], \\g_{\omega\eta\gamma} &= g\left\{\left[\left(\frac{Z_{\text{NS}}}{3} - \epsilon_{12}\right)c\phi_{23} + \epsilon_{13}s\phi_{23}\right]c\phi_V\right. \\&\quad \left.- \frac{2}{3}z_S\frac{\bar{m}}{m_s}s\phi_{23}s\phi_V\right\}, \\g_{\eta'\omega\gamma} &= g\left\{\left[\left(\frac{Z_{\text{NS}}}{3} - \epsilon_{12}\right)s\phi_{23} - \epsilon_{13}c\phi_{23}\right]c\phi_V\right. \\&\quad \left.+ \frac{2}{3}z_S\frac{\bar{m}}{m_s}c\phi_{23}s\phi_V\right\}, \\g_{\phi\pi^0\gamma} &= g\left[\left(1 + \frac{\epsilon_{12}}{3}Z_{\text{NS}}\right)s\phi_V - \frac{2}{3}z_S\frac{\bar{m}}{m_s}\epsilon_{13}c\phi_V\right],\end{aligned}$$

$$\begin{aligned}g_{\phi\pi^0\gamma} &= g\left[\left(1 + \frac{\epsilon_{12}}{3}Z_{\text{NS}}\right)s\phi_V - \frac{2}{3}z_S\frac{\bar{m}}{m_s}\epsilon_{13}c\phi_V\right], \\g_{\phi\eta\gamma} &= g\left\{\left[\left(\frac{Z_{\text{NS}}}{3} - \epsilon_{12}\right)c\phi_{23} + \epsilon_{13}s\phi_{23}\right]s\phi_V\right. \\&\quad \left.+ \frac{2}{3}z_S\frac{\bar{m}}{m_s}s\phi_{23}c\phi_V\right\}, \\g_{\phi\eta'\gamma} &= g\left\{\left[\left(\frac{Z_{\text{NS}}}{3} - \epsilon_{12}\right)s\phi_{23} - \epsilon_{13}c\phi_{23}\right]s\phi_V\right. \\&\quad \left.- \frac{2}{3}z_S\frac{\bar{m}}{m_s}c\phi_{23}c\phi_V\right\}, \\g_{K^{*0}K^0\gamma} &= -\frac{1}{3}g\left(1 + \frac{\bar{m}}{m_s}\right)z_{K^0} = -\frac{1}{3}g\left(1 + z_S\frac{\bar{m}}{m_s}\right)z'_{K^0}, \\g_{K^{*+}K^+\gamma} &= \frac{1}{3}g\left(2 - \frac{\bar{m}}{m_s}\right)z_{K^+} = \frac{1}{3}g\left(2 - z_S\frac{\bar{m}}{m_s}\right)z'_{K^+},\end{aligned}$$

$$\begin{aligned}z_{\text{NS}} &= Z_{\text{NS}}/Z_3 & z_S &= Z_S/Z_3 & z_+ &= Z_+/Z_3 \\z_{K^0} &= Z_{K^0}/Z_3 & z_{K^+} &= Z_{K^+}/Z_3 & g &= Z_3 g_e\end{aligned}$$

Results

η - η' mixing revisited

Table 1

Comparison between estimations for the seven free parameters from the model presented in Ref. [6], using the PDG 2000 and the most up-to-date experimental data.

| Parameter | Estimation from [6] | Current Estimation |
|-------------------------------|----------------------------------|----------------------------------|
| g | $0.70 \pm 0.02 \text{ GeV}^{-1}$ | $0.70 \pm 0.01 \text{ GeV}^{-1}$ |
| $\frac{m_s}{\overline{m}}$ | 1.24 ± 0.07 | 1.17 ± 0.06 |
| ϕ_P | $(37.7 \pm 2.4)^\circ$ | $(41.4 \pm 0.5)^\circ$ |
| ϕ_V | $(3.4 \pm 0.2)^\circ$ | $(3.3 \pm 0.1)^\circ$ |
| z_{NS} | 0.91 ± 0.05 | 0.84 ± 0.02 |
| z_S | 0.89 ± 0.07 | 0.76 ± 0.04 |
| z_K | 0.91 ± 0.04 | 0.89 ± 0.03 |
| $\chi^2_{\min}/\text{d.o.f.}$ | 0.7 | 4.6 |

Results


FIT 1

$$\begin{aligned} g &= 0.69 \pm 0.01 \text{ GeV}^{-1}, & z_+ &= 0.95 \pm 0.05, \\ \phi_{23} &= (41.5 \pm 0.5)^\circ, & \phi_V &= (4.0 \pm 0.2)^\circ, \\ \epsilon_{12} &= (2.3 \pm 1.0)\%, & \epsilon_{13} &= (2.5 \pm 0.9)\%, \\ z_{\text{NS}} &= 0.89 \pm 0.03, & z_S \bar{m}/m_S &= 0.65 \pm 0.01, \\ z'_{K^0} &= 1.01 \pm 0.04, & z'_{K^+} &= 0.76 \pm 0.04. \end{aligned}$$

$$\chi^2_{\min}/\text{dof} \simeq 4.6/2 = 2.3$$

ϕ_{23}, ϕ_V **very good agreement** with recent published results

$\epsilon_{12}, \epsilon_{13}$ **very small but not compatible with zero,**
with a CL of 2.3 and 2.8 sigmas, respectively



$$\epsilon = \epsilon_{\pi\eta} = (0.1 \pm 0.9)\%$$

$$\epsilon' = \epsilon_{\pi\eta'} = (3.4 \pm 0.9)\% \quad \textbf{CL 3.8 sig.}$$

Results

FIT 2

Turning off **secondary mechanism** of **isospin breaking**

$$z_+ = 1 \text{ \& } z_{K^0} = z_K^+$$

$$g = 0.69 \pm 0.01 \text{ GeV}^{-1}, m_s/\bar{m} = 1.17 \pm 0.06,$$

$$\phi_{23} = (41.5 \pm 0.5)^\circ, \quad \phi_V = (4.0 \pm 0.2)^\circ,$$


$$\epsilon_{12} = (2.4 \pm 1.0)\%, \quad \epsilon_{13} = (2.5 \pm 0.9)\%,$$

$$z_{NS} = 0.89 \pm 0.03, \quad z_S = 0.77 \pm 0.04,$$

$$z_K = 0.90 \pm 0.03,$$

$$\chi_{\min}^2/\text{dof} \simeq 5.6/3 = 1.9$$

The z's are still different from zero



Secondary mechanism of **SU(3)_F breaking** is still required

Results

FIT 3

Using Kroll's results: $\epsilon_{12} = (1.6 \pm 0.2)\%$ $\epsilon_{13} = (-0.8 \pm 0.1)\%$

$$\begin{aligned} g &= 0.69 \pm 0.01 \text{ GeV}^{-1}, m_s/\bar{m} = 1.17 \pm 0.06, \\ \phi_{23} &= (41.4 \pm 0.5)^\circ, \quad \phi_V = (3.1 \pm 0.1)^\circ, \\ z_{NS} &= 0.86 \pm 0.0, \quad z_S = 0.77 \pm 0.04, \\ z_K &= 0.90 \pm 0.03, \end{aligned}$$

$$\chi^2_{\min}/\text{dof} \simeq 22.0/5 = 4.4$$

Results

FIT 4

Using Escribano et al's results:

$$\epsilon_{12} = (7.5 \pm 0.2) \times 10^{-3} \quad \epsilon_{13} = (-6.3 \pm 0.2) \times 10^{-3}$$


$$g = 0.70 \pm 0.01 \text{ GeV}^{-1}, m_s/\bar{m} = 1.17 \pm 0.06,$$

$$\phi_{23} = (41.4 \pm 0.5)^\circ, \quad \phi_V = (3.2 \pm 0.1)^\circ,$$

$$z_{NS} = 0.85 \pm 0.02, \quad z_S = 0.77 \pm 0.04,$$

$$z_K = 0.90 \pm 0.03,$$

$$\chi^2_{\min}/\text{dof} \simeq 24.0/5 = 4.8$$



Theoretical estimations by Kroll and Escribano et al.

do not appear to agree with the **most recent experimental data**

Results

FIT 5

Charged and neutral **kaon transitions** are **not considered**

$$\begin{aligned} g &= 0.69 \pm 0.01 \text{ GeV}^{-1}, z_S \bar{m}/m_S = 0.65 \pm 0.01, \\ \phi_{23} &= (41.5 \pm 0.5)^\circ, \quad \phi_V = (4.0 \pm 0.2)^\circ, \\ \epsilon_{12} &= (2.4 \pm 1.0)\%, \quad \epsilon_{13} = (2.5 \pm 0.9)\%, \\ z_{NS} &= 0.89 \pm 0.03. \end{aligned}$$

$$\chi^2_{\min}/\text{dof} \simeq 5.6/3 = 1.9$$

$\epsilon_{12}, \epsilon_{13}$ again **incompatible with zero**,
with a CL of 2.4 and 2.8 sigmas, respectively

Results

Summary of Fits

Table 2

Summary of fitted values for the Fit 1, Fit 2, Fit 3, Fit 4 and Fit 5, corresponding to Eqs. (13), (14), (15), (16), and (17), respectively.


| Parameter | Fit 1 | Fit 2 | Fit 3 | Fit 4 | Fit 5 |
|-------------------------------------|-------------------|-------------------|-----------------|-----------------|-------------------|
| $g \text{ (GeV}^{-1}\text{)}$ | 0.69 ± 0.01 | 0.69 ± 0.01 | 0.69 ± 0.01 | 0.70 ± 0.01 | 0.69 ± 0.01 |
| ϵ_{12} | $(2.3 \pm 1.0)\%$ | $(2.4 \pm 1.0)\%$ | - | - | $(2.4 \pm 1.0)\%$ |
| ϵ_{13} | $(2.5 \pm 0.9)\%$ | $(2.5 \pm 0.9)\%$ | - | - | $(2.5 \pm 0.9)\%$ |
| $\phi_{23} \text{ (}^\circ\text{)}$ | 41.5 ± 0.5 | 41.5 ± 0.05 | 41.4 ± 0.5 | 41.4 ± 0.5 | 41.5 ± 0.5 |
| $\phi_V \text{ (}^\circ\text{)}$ | 4.0 ± 0.2 | 4.0 ± 0.2 | 3.1 ± 0.1 | 3.2 ± 0.1 | 4.0 ± 0.2 |
| m_s/\overline{m} | - | 1.17 ± 0.06 | 1.17 ± 0.06 | 1.17 ± 0.06 | - |
| $z_S \overline{m}/m_s$ | 0.65 ± 0.01 | - | - | - | 0.65 ± 0.01 |
| z_{NS} | 0.89 ± 0.03 | 0.89 ± 0.03 | 0.86 ± 0.02 | 0.85 ± 0.02 | 0.89 ± 0.03 |
| z_+ | 0.95 ± 0.05 | - | - | - | - |
| z_S | - | 0.77 ± 0.04 | 0.77 ± 0.04 | 0.77 ± 0.04 | - |
| z_K | - | 0.90 ± 0.03 | 0.90 ± 0.03 | 0.90 ± 0.03 | - |
| z'_{K^0} | 1.01 ± 0.04 | - | - | - | - |
| z'_{K^+} | 0.76 ± 0.04 | - | - | - | - |
| $\chi^2_{\min}/\text{d.o.f.}$ | 2.3 | 1.9 | 4.4 | 4.8 | 1.9 |

Conclusions

- The quality of the most **up-to-date experimental data** enables a small amount of **isospin-symmetry breaking** that is inconsistent with zero with a CL of aprox. 2.5 sigmas
- The quality of the performed fits is **good**
- The **estimations** for the fit parameters appear to be robust
- Our **estimates** for

$$\epsilon_{12} = (2.4 \pm 1.0)\%$$

$$\epsilon_{13} = (2.5 \pm 0.9)\%$$



$$\epsilon = \epsilon_{\pi\eta} = (0.1 \pm 0.9)\%$$

$$\epsilon' = \epsilon_{\pi\eta'} = (3.5 \pm 0.9)\%$$

are not in accordance with **theoretical estimates**