

# Hadronic contribution to $(g - 2)_\mu$ in the Standard Model: data-driven approach

Gilberto Colangelo

*u*<sup>b</sup>

---

<sup>b</sup>  
UNIVERSITÄT  
BERN

AEC  
ALBERT EINSTEIN CENTER  
FOR FUNDAMENTAL PHYSICS

vQCHS Conference 2021  
Online, August 2-6, 2021

# Outline

Introduction:  $(g - 2)_\mu$  in the Standard Model


Hadronic Vacuum Polarization contribution to  $(g - 2)_\mu$

Hadronic light-by-light contribution to  $(g - 2)_\mu$

Conclusions and Outlook


# Present status of $(g - 2)_\mu$ , experiment vs SM

Before the Fermilab result



# Present status of $(g-2)_\mu$ , experiment vs SM

After the Fermilab result



# Present status of $(g - 2)_\mu$ , experiment vs SM

$$a_\mu(BNL) = 116\,592\,089(63) \times 10^{-11}$$

$$a_\mu(FNAL) = 116\,592\,040(54) \times 10^{-11}$$

$$a_\mu(Exp) = 116\,592\,061(41) \times 10^{-11}$$

→ talk by E. Valetov

# White Paper (2020): $(g - 2)_\mu$ , experiment vs SM

Contribution	Value $\times 10^{11}$
HVP LO ( $e^+e^-$ )	6931(40)
HVP NLO ( $e^+e^-$ )	-98.3(7)
HVP NNLO ( $e^+e^-$ )	12.4(1)
HVP LO (lattice, $udsc$ )	7116(184)
HLbL (phenomenology)	92(19)
HLbL NLO (phenomenology)	2(1)
HLbL (lattice, $uds$ )	79(35)
HLbL (phenomenology + lattice)	90(17)
QED	116 584 718.931(104)
Electroweak	153.6(1.0)
HVP ( $e^+e^-$ , LO + NLO + NNLO)	6845(40)
HLbL (phenomenology + lattice + NLO)	92(18)
Total SM Value	116 591 810(43)
Experiment	116 592 061(41)
Difference: $\Delta a_\mu := a_\mu^{\text{exp}} - a_\mu^{\text{SM}}$	251(59)

White Paper (2020):  $(g - 2)_\mu$ , experiment vs SM

Contribution	Value $\times 10^{11}$
HVP LO ( $e^+e^-$ )	6931(40)
HVP NLO ( $e^+e^-$ )	-98.3(7)
HVP NNLO ( $e^+e^-$ )	12.4(1)
HVP LO (lattice <b>BMW(20)</b> , $udsc$ )	<b>7075(55)</b>
HLbL (phenomenology)	92(19)
HLbL NLO (phenomenology)	2(1)
HLbL (lattice, $uds$ )	79(35)
HLbL (phenomenology + lattice)	90(17)
QED	116 584 718.931(104)
Electroweak	153.6(1.0)
HVP ( $e^+e^-$ , LO + NLO + NNLO)	6845(40)
HLbL (phenomenology + lattice + NLO)	92(18)
Total SM Value	116 591 810(43)
Experiment	<b>116 592 061(41)</b>
Difference: $\Delta a_\mu := a_\mu^{\text{exp}} - a_\mu^{\text{SM}}$	251(59)

# White Paper (2020): $(g - 2)_\mu$ , experiment vs SM

## White Paper:

T. Aoyama et al. Phys. Rep. 887 (2020) = WP(20)

## Muon $g - 2$ Theory Initiative

### Steering Committee:

GC

Michel Davier (vice-chair)

Simon Eidelman

Aida El-Khadra (chair)

Martin Hoferichter

Christoph Lehner (vice-chair)

Tsutomu Mibe (J-PARC E34 experiment)

(Andreas Nyffeler until summer 2020)

Lee Roberts (Fermilab E989 experiment)

Thomas Teubner

Hartmut Wittig



# White Paper (2020): $(g - 2)_\mu$ , experiment vs SM

## White Paper:

T. Aoyama et al. Phys. Rep. 887 (2020) = WP(20)



## Muon $g - 2$ Theory Initiative

### Workshops:

- First plenary meeting, Q-Center (Fermilab), 3-6 June 2017
- HVP WG workshop, KEK (Japan), 12-14 February 2018
- HLbL WG workshop, U. of Connecticut, 12-14 March 2018
- Second plenary meeting, Mainz, 18-22 June 2018
- Third plenary meeting, Seattle, 9-13 September 2019
- Lattice HVP workshop, virtual, 16-20 November 2020
- Fourth plenary meeting, KEK (virtual), 28 June-02 July 2021

# Theory uncertainty comes from hadronic physics

- ▶ Hadronic contributions responsible for most of the theory uncertainty → see also talk by M. Della Morte
- ▶ Hadronic vacuum polarization (HVP) is  $\mathcal{O}(\alpha^2)$ , dominates the total uncertainty, despite being known to  $< 1\%$




- ▶ unitarity and analyticity  $\Rightarrow$  dispersive approach
- ▶  $\Rightarrow$  direct relation to experiment:  $\sigma_{\text{tot}}(e^+e^- \rightarrow \text{hadrons})$
- ▶  $e^+e^-$  Exps: BaBar, Belle, BESIII, CMD2/3, KLOE2, SND
- ▶ **alternative approach**: lattice, becoming competitive

(BMW, ETMC, Fermilab, HPQCD, Mainz, MILC, RBC/UKQCD)

→ talk by Z. Fodor

# Theory uncertainty comes from hadronic physics

- ▶ Hadronic contributions responsible for most of the theory uncertainty → see also talk by M. Della Morte
- ▶ Hadronic vacuum polarization (HVP) is  $\mathcal{O}(\alpha^2)$ , dominates the total uncertainty, despite being known to  $< 1\%$
- ▶ Hadronic light-by-light (HLbL) is  $\mathcal{O}(\alpha^3)$ , known to  $\sim 20\%$ , second largest uncertainty (now subdominant)




- ▶ 4-point fct. of em currents in QCD: more complicated than HVP
- ▶ **recently**: dispersive approach  $\Rightarrow$  data-driven, systematic treatment
- ▶ lattice QCD is becoming competitive

(Mainz, RBC/UKQCD)

# HVP contribution: Master Formula

Unitarity relation: **simple**, same for all intermediate states



$$\text{Im}\Pi(q^2) \propto \sigma(e^+e^- \rightarrow \text{hadrons}) = \sigma(e^+e^- \rightarrow \mu^+\mu^-)R(q^2)$$

Analyticity  $\Rightarrow$  **Master formula for HVP**

Bouchiat, Michel (61)

$$a_\mu^{\text{hvp}} = \frac{\alpha^2}{3\pi^2} \int_{s_{th}}^{\infty} \frac{ds}{s} K(s) R(s)$$

$K(s)$  known, depends on  $m_\mu$  and  $K(s) \sim \frac{1}{s}$  for large  $s$

# Comparison between DHMZ19 and KNT19

	DHMZ19	KNT19	Difference
$\pi^+\pi^-$	507.85(0.83)(3.23)(0.55)	504.23(1.90)	3.62
$\pi^+\pi^-\pi^0$	46.21(0.40)(1.10)(0.86)	46.63(94)	-0.42
$\pi^+\pi^-\pi^+\pi^-$	13.68(0.03)(0.27)(0.14)	13.99(19)	-0.31
$\pi^+\pi^-\pi^0\pi^0$	18.03(0.06)(0.48)(0.26)	18.15(74)	-0.12
$K^+K^-$	23.08(0.20)(0.33)(0.21)	23.00(22)	0.08
$K_S K_L$	12.82(0.06)(0.18)(0.15)	13.04(19)	-0.22
$\pi^0\gamma$	4.41(0.06)(0.04)(0.07)	4.58(10)	-0.17
Sum of the above	626.08(0.95)(3.48)(1.47)	623.62(2.27)	2.46
[1.8, 3.7] GeV (without $c\bar{c}$ )	33.45(71)	34.45(56)	-1.00
$J/\psi, \psi(2S)$	7.76(12)	7.84(19)	-0.08
[3.7, $\infty$ ) GeV	17.15(31)	16.95(19)	0.20
Total $a_\mu^{\text{HVP, LO}}$	694.0(1.0)(3.5)(1.6)(0.1) $_{\psi(0.7)\text{DV+QCD}}$	692.8(2.4)	1.2

DHMZ = Davier, Hoecker, Malaescu, Zhang,

KNT = Keshavarzi, Nomura, Teubner

## $2\pi$ : comparison with the dispersive approach

The  $2\pi$  channel can itself be described dispersively  $\Rightarrow$  more constrained theoretically

Ananthanarayan, Caprini, Das (19), GC, Hoferichter, Stoffer (18)

Energy range	ACD18	CHS18	DHMZ19	KNT19
$< 0.6$ GeV		110.1(9)	110.4(4)(5)	108.7(9)
$< 0.7$ GeV		214.8(1.7)	214.7(0.8)(1.1)	213.1(1.2)
$< 0.8$ GeV		413.2(2.3)	414.4(1.5)(2.3)	412.0(1.7)
$< 0.9$ GeV		479.8(2.6)	481.9(1.8)(2.9)	478.5(1.8)
$\leq 1.0$ GeV		495.0(2.6)	497.4(1.8)(3.1)	493.8(1.9)
[0.6, 0.7] GeV		104.7(7)	104.2(5)(5)	104.4(5)
[0.7, 0.8] GeV		198.3(9)	199.8(0.9)(1.2)	198.9(7)
[0.8, 0.9] GeV		66.6(4)	67.5(4)(6)	66.6(3)
[0.9, 1.0] GeV		15.3(1)	15.5(1)(2)	15.3(1)
$\leq 0.63$ GeV	132.9(8)	132.8(1.1)	132.9(5)(6)	131.2(1.0)
[0.6, 0.9] GeV		369.6(1.7)	371.5(1.5)(2.3)	369.8(1.3)
$[\sqrt{0.1}, \sqrt{0.95}]$ GeV		490.7(2.6)	493.1(1.8)(3.1)	489.5(1.9)

## Combination method and final result

Complete analyses DHMZ19 and KNT19, as well as CHS19 ( $2\pi$ ) and HHK19 ( $3\pi$ ), have been so combined:


- ▶ central values are obtained by simple averages (for each channel and mass range)
- ▶ the largest experimental and systematic uncertainty of DHMZ and KNT is taken
- ▶ 1/2 difference DHMZ–KNT (or BABAR–KLOE in the  $2\pi$  channel, if larger) is added to the uncertainty

**Final result:**

$$\begin{aligned}
 a_\mu^{\text{HVP, LO}} &= 693.1(2.8)_{\text{exp}}(2.8)_{\text{sys}}(0.7)_{\text{DV+QCD}} \times 10^{-10} \\
 &= 693.1(4.0) \times 10^{-10}
 \end{aligned}$$

# What if the BMW result is right?

→ talk by Z. Fodor





# Consequences of the BMW result

A shift in the value of  $a_\mu^{\text{HVP, LO}}$  would have consequences:

- ▶  $\Delta a_\mu^{\text{HVP, LO}} \Leftrightarrow \Delta \sigma(e^+ e^- \rightarrow \text{hadrons})$
- ▶  $\Delta \alpha_{\text{had}}(M_Z^2)$  is determined by an integral of the same  $\sigma(e^+ e^- \rightarrow \text{hadrons})$  (more weight at high energy)
- ▶ changing  $a_\mu^{\text{HVP, LO}}$  necessarily implies a shift in  $\Delta \alpha_{\text{had}}(M_Z^2)$ :  
 $\Rightarrow$  impact on the EW-fit
- ▶ to save the EW-fit  $\Delta \sigma(e^+ e^- \rightarrow \text{hadrons})$  must occur below  $\sim 1$  (max 2) GeV

Crivellin, Hoferichter, Manzari, Montull (20)/Keshavarzi, Marciano, Passera, Sirlin (20)/Malaescu, Schott (20)


- ▶ or the need for BSM physics would be moved elsewhere

for a BSM perspective  $\rightarrow$  talk by A. Crivellin

# Changes in $\sigma(e^+e^- \rightarrow \text{hadrons})$ below 1 GeV?

- ▶ Below 1 – 2 GeV only one significant channel:  $\pi^+\pi^-$
- ▶ Strongly constrained by analyticity and unitarity ( $F_\pi^V(s)$ )
- ▶  $F_\pi^V(s)$  parametrization which satisfies these  
 $\Rightarrow$  small number of parameters GC, Hoferichter, Stoffer (18)
- ▶  $\Delta a_\mu^{\text{HVP, LO}} \Leftrightarrow$  shifts in these parameters  
analysis of the corresponding scenarios GC, Hoferichter, Stoffer (21)

# Changes in $\sigma(e^+e^- \rightarrow \text{hadrons})$ below 1 GeV?




$c_k \Leftrightarrow$  inelastic channels

GC, Hoferichter, Stoffer (21)

Tension [BMW20 vs  $e^+e^-$  data] stronger for KLOE than for BABAR

# Changes in $\sigma(e^+e^- \rightarrow \text{hadrons})$ below 1 GeV?





$c_k \Leftrightarrow$  inelastic channels

GC, Hoferichter, Stoffer (21)

$$10^4 \Delta\alpha_{\text{had}}^{(5)}(M_Z^2) = \begin{cases} 272.2(4.1) & \text{EW fit} \\ 276.1(1.1) & \sigma_{\text{had}}(s) \end{cases}$$


# BMW vs individual $\pi^+\pi^-$ experiments



BMW vs individual  $\pi^+\pi^-$  experiments


BMW vs individual  $\pi^+\pi^-$  experiments

## Article




# BMW vs individual $\pi^+\pi^-$ experiments

## Weight functions for the window quantities





# BMW vs individual $\pi^+\pi^-$ experiments




$a_\mu^{\text{win}}$  suggests that  $\sim 5 \times 10^{-10}$  must come from above 1 GeV

# Calculating the HLbL contribution

Calculating the HLbL contribution is complicated → talk by M. Hoferichter

- ▶ 4-point function of em currents in QCD



- ▶ a data-driven approach like for HVP seemed hopeless but has been recently developed and used

GC, Hoferichter, Procura, Stoffer=CHPS (14,15,17), Hoferichter, Hoid, Kubis, Leupold, Schneider (18)

- ▶ lattice QCD is an alternative and is making fast progress

RBC/UKQCD (20), Mainz (19,20)

# HLbL contribution: Master Formula

$$a_\mu^{\text{HLbL}} = \frac{2\alpha^3}{48\pi^2} \int_0^\infty dQ_1 \int_0^\infty dQ_2 \int_{-1}^1 d\tau \sqrt{1 - \tau^2} \sum_{i=1}^{12} T_i(Q_1, Q_2, \tau) \bar{\Pi}_i(Q_1, Q_2, \tau)$$


$Q_i^\mu$  are the **Wick-rotated** four-momenta and  $\tau$  the four-dimensional angle between Euclidean momenta:

$$Q_1 \cdot Q_2 = |Q_1| |Q_2| \tau$$

The integration variables  $Q_1 := |Q_1|$ ,  $Q_2 := |Q_2|$ .

CHPS (15)

- ▶  $T_i$ : known kernel functions
- ▶  $\bar{\Pi}_i$  are amenable to a dispersive treatment: **their imaginary parts are related to measurable subprocesses**



# Improvements obtained with the dispersive approach

Contribution	PdRV(09) <i>Glasgow consensus</i>	N/JN(09)	J(17)	WP(20)
$\pi^0, \eta, \eta'$ -poles	114(13)	99(16)	95.45(12.40)	93.8(4.0)
$\pi, K$ -loops/boxes	-19(19)	-19(13)	-20(5)	-16.4(2)
$S$ -wave $\pi\pi$ rescattering	-7(7)	-7(2)	-5.98(1.20)	-8(1)
subtotal	88(24)	73(21)	69.5(13.4)	69.4(4.1)
scalars	—	—	—	} - 1(3)
tensors	—	—	1.1(1)	
axial vectors	15(10)	22(5)	7.55(2.71)	
$u, d, s$ -loops / short-distance	—	21(3)	20(4)	6(6)
				15(10)
$c$ -loop	2.3	—	2.3(2)	3(1)
total	105(26)	116(39)	100.4(28.2)	92(19)

- ▶ significant reduction of uncertainties in the first three rows:  
low-energy region well constrained by a dispersive approach


CHPS (17), Masjuan, Sánchez-Puertas (17) Hoferichter, Hoid et al. (18), Gerardin, Meyer, Nyffeler (19)

- ▶ 1 – 2 GeV and asymptotic region (short distance constraints)  
have been improved, but still work in progress (see WP(20))

Melnikov, Vainshtein (04), (.....), Bijnsens, Hermansson-Truedsson, Laub, Rodríguez-Sánchez (20,21)

→ talks by Hoferichter and Rodríguez-Sánchez

# Situation for HLbL



# Conclusions

- ▶ The WP provides the current status of the SM evaluation of  $(g - 2)_\mu$ :  $4.2\sigma$  **discrepancy with experiment (w/ FNAL)**
- ▶ Evaluation of the HVP contribution based on the dispersive approach: **0.6% error**  $\Rightarrow$  **dominates the theory uncertainty**
- ▶ Recent lattice calculation [BMW(20)] has reached a similar precision but **differs from the dispersive one** (=from  $e^+e^-$  data).  
If confirmed  $\Rightarrow$  discrepancy with experiment  $\searrow$  **below  $2\sigma$**
- ▶ Evaluation of the HLbL contribution based on the dispersive approach: **20% accuracy**. Two recent lattice calculations [RBC/UKQCD(20), Mainz(21)] agree with it

# Outlook

- ▶ The Fermilab experiment aims to reduce the BNL uncertainty by a **factor four**  $\Rightarrow$  potential  **$7\sigma$**  discrepancy
- ▶ Improvements on the SM theory side:
  - ▶ HVP data-driven:  
Other  $e^+e^-$  experiments are available or forthcoming:  
**SND, BaBar, Belle II, BESIII, CMD3**  $\Rightarrow$  **Error reduction**  
**MuonE** will provide an alternative way to measure HVP
  - ▶ HVP lattice:  
**BMW** result must be confirmed (or refuted) by others.  
**Difference to data-driven evaluation must be understood**
  - ▶ HLbL data-driven: goal of  **$\sim 10\%$  uncertainty** within reach
  - ▶ HLbL lattice: **RBC/UKQCD**  $\Rightarrow$  similar precision as **Mainz**.  
**Good agreement with data-driven evaluation.**

# Future: Muon $g - 2$ /EDM experiment @ J-PARC

