

# QCD vacuum replicas are metastable

talk at XVth Quark Confinement and the Hadron Spectrum, August 1- 7, Stavanger

Pedro Bicudo (speaker) and J.Emilio Ribeiro

CeFEMA, Departamento de Física,  
Instituto Superior Técnico  
Universidade de Lisboa, Portugal

*bicudo@tecnico.ulisboa.pt*

August 05, 2022

## Abstract

We study the meson spectra in the excited QCD vacua, denominated replicas. We find all mesons have real masses, with no tachyons, thus showing the QCD replicas are indeed metastable.

## Introduction

- Spontaneous breaking of chiral symmetry and the QCD vacua
- Approach of Gaussian approximation to QCD with two cumulants
- Our previous knowledge of the replicas

## Details of the mass gap and bound state equations

- Mass gap equation and replicas
- Salpeter-RPA equations for mesons
- Numerical solution of the mass gap and boundstate equations.

## Summary

## Introduction

Spontaneous breaking of chiral symmetry and the QCD vacua

Approach of Gaussian approximation to QCD with two cumulants

Our previous knowledge of the replicas

## Details of the mass gap and bound state equations

Mass gap equation and replicas

Salpeter-RPA equations for mesons

Numerical solution of the mass gap and boundstate equations.

## Summary

In 1960 Gell-Mann and Levy, with the sigma model of Fig. 3, and soon after Nambu and Jona-Lasinio modeled for the first time the spontaneous breaking of chiral symmetry.




Figure: The sigma model is 62 years old, what have we learned since then?

This is an effective  $\phi^4$  model:

1. there are two extrema,
2. the particle masses<sup>2</sup> are the vertical curvature,
3. in the true vacuum,  $m_\sigma^2 > 0$ ,  $m_\pi^2 = 0$
4. in the false vacuum,  $m_\sigma^2 < 0$ ,  $m_\pi^2 < 0$ , we have tachyons : unstable vacuum
5. it is not renormalizable, not eligible as a field theory

The sigma model advanced physics a lot and is still used in Higgs sector of the standard model. However it is just a model, it is not renormalizable.

In particle physics we have a theory with spontaneous breaking of chiral symmetry: QCD., as verified by lattice QCD simulations. However we don't yet know how to study the QCD vacuum with the lattice.

In this talk I will show theoretical evidence that the QCD vacuum has a much richer structure than the sigma model, with a tower of metastable vacua. We denominate these novel vacua the replicas of QCD.




Figure: This are the extrema we expect for QCD.

## Introduction

Spontaneous breaking of chiral symmetry and the QCD vacua

**Approach of Gaussian approximation to QCD with two cumulants**

Our previous knowledge of the replicas

## Details of the mass gap and bound state equations

Mass gap equation and replicas



Salpeter-RPA equations for mesons

Numerical solution of the mass gap and boundstate equations.

## Summary



mass gap equation

stationary vacuum  = 0 or energy density extremum  = 0 (i)

Bethe-Salpeter equation, for bound state mass and vertex

 =  also for baryons... (ii)

Note: (i) and (ii) are formally the same when the meson  $M$  is the pseudoscalar  $\pi$  with zero mass


All the other hadrons get a finite mass: 38% of Nucleon mass

Figure: Examples of what the two cumulant approach implies.

Importantly, all the PCAC theorems, such as the Gell-Mann Oakes and Renner relation for  $M_\pi$  and  $\langle \bar{\psi}\psi \rangle$ , the Goldberger and Treiman relation and the Weinberg Theorem for  $\pi\pi$  scattering, the Adler zero for pion couplings are verified [17, 20, 9].

Moreover this generates a constituent mass for the light quark, leading to chiral symmetry restoration for excited hadrons, while for heavy quarks, a quasi classical regime for hadrons must set in [15].

In Eq. (1),  $m_0$  is the current mass of the quark, for simplicity we will set  $m_0$  to zero.



**Figure:** The dependence [3] of the constituent quark mass  $m_c(k)$  on the current quark mass  $m_0$ .

The big picture is independent of the details of the kernel: all the results will apply equally well with any confining potential. In this talk and for numerical simplicity, we will use the harmonic confining kernel.

The model of Eq.(1) was suggested in the mid-eighties [1, 17, 18], and latter re-derived in terms of coherent states of  $^3P_0$  quark-antiquark pairs [11, 12].


## Introduction

Spontaneous breaking of chiral symmetry and the QCD vacua  
Approach of Gaussian approximation to QCD with two cumulants  
**Our previous knowledge of the replicas**

## Details of the mass gap and bound state equations

Mass gap equation and replicas  
Salpeter-RPA equations for mesons  
Numerical solution of the mass gap and boundstate equations.

## Summary




**Figure:** The first solution,  $\varphi_0(k)$ , of the mass gap equation, representing the vacuum  $|0\rangle$  and its two first two other solution,  $\varphi(k)_1$  and  $\varphi(k)_2$  for a linear confining potential [7, 4], corresponding to the truncated Coulomb gauge of QCD), computed in the chiral limit of  $m_q \rightarrow 0$ .

With confining kernels, there is an infinite tower of excited vacua-like states, the replicas  $|\mathcal{R}1\rangle$ , interpolating between the true chiral symmetry breaking, physical vacuum, to the highest chiral invariant vacuum [17, 11, 7]. Examples of solutions in the case of the linear potential [7] for the chiral angle  $\varphi$  are depicted in Fig. 6.

The mass gap equation for the vacua has similarities with the Bethe Salpeter equation for Hadrons in the true vacuum, who also have an infinite tower of excited states. This is a robust evidence for the replicas.

We also know that the number of tachyon states for the false chiral invariant vacuum becomes infinite [3] in the pseudoscalar and scalar channels, whereas, from the true vacuum, we have an infinite set of true hadronic states. This shows the false vacuum is unstable while the true vacuum  $|0\rangle$  is stable, it is an absolute minimum.

For infinite volumes, the different vacua correspond to orthogonal  ${}^3P_0$  coherent states so that, two different coherent states, have a zero overlap between them, hence being independent of each other. In addition, they are separated apart by an infinite amount of energy [8, 2]. For a finite volume, see Fig. 7.



**Figure:** We show (blue) the overlap  $\langle \mathcal{R}1|0\rangle$  and (orange) the corresponding energy  $E_{\mathcal{R}1}$  of the first replica for different spherical bubble radii. The energy density is ( $\mathcal{E}_{\mathcal{R}1} \simeq 87.2476 \text{ MeV}/\text{fm}^3$  (using  $K_0 \simeq 300 \text{ MeV}$ ) and, for instance, with finite a radius of only 5 fm, we would had to add to any hadronic excitation in  $\mathcal{R}1$  another 45 GeV, and this for an overlap  $\langle \mathcal{R}1|0\rangle = 2.9 \times 10^{-16}$ .

## Introduction

Spontaneous breaking of chiral symmetry and the QCD vacua  
Approach of Gaussian approximation to QCD with two cumulants  
Our previous knowledge of the replicas

## Details of the mass gap and bound state equations

**Mass gap equation and replicas**

Salpeter-RPA equations for mesons

Numerical solution of the mass gap and boundstate equations.

## Summary

There are three equivalent methods to solve the mass gap equation:

1. assume a quark-antiquark  ${}^3P_0$  coherent state, and minimize its correspondent energy density,
2. rotate the quark fields by a Bogoliubov-Valatin canonical transformation to diagonalize the Hamiltonian,
3. solve the Schwinger-Dyson equations for the quark propagators.

Any of these methods lead to the same mass gap equation and  $m_c(k)$ .

Technically, it is convenient to decompose the relativistic invariant Dirac-Feynman propagators [17, 10], in the quark and antiquark Bethe-Goldstone propagators, with quark and antiquark spinors  $u_s$  and  $v_s$ , close to the formalism of non-relativistic quark models,

$$S_{Dirac}(k_0, \vec{k}) = \frac{i}{\not{k} - m + i\epsilon} = \frac{\sum_s u_s u_s^\dagger \beta}{k_0 - E(k) + i\epsilon} - \frac{\sum_s v_s v_s^\dagger \beta}{-k_0 - E(k) + i\epsilon} \quad (3)$$

Replacing the propagator of Eq. (3) in its Schwinger-Dyson equation,

$$\begin{aligned} 0 &= u_s^\dagger(k) \left\{ k\hat{k} \cdot \vec{\alpha} + m_0\beta - \int \frac{dw'}{2\pi} \frac{d^3k'}{(2\pi)^3} iV(k-k') \right. \\ &\quad \left. \sum_{s'} \left[ \frac{u(k')_{s'} u^\dagger(k')_{s'}}{w' - E(k') + i\epsilon} - \frac{v(k')_{s'} v^\dagger(k')_{s'}}{-w' - E(k') + i\epsilon} \right] \right\} v_{s'}(k) \\ E(k) &= u_s^\dagger(k) \left\{ k\hat{k} \cdot \vec{\alpha} + m_0\beta - \int \frac{dw'}{2\pi} \frac{d^3k'}{(2\pi)^3} iV(k-k') \right. \\ &\quad \left. \sum_{s'} \left[ \frac{u(k')_{s'} u^\dagger(k')_{s'}}{w' - E(k') + i\epsilon} - \frac{v(k')_{s'} v^\dagger(k')_{s'}}{-w' - E(k') + i\epsilon} \right] \right\} u_s(k), \end{aligned} \quad (4)$$

It was shown that for the wide class of confining potentials [7], with the exponent  $0 < \alpha \leq 2$  – see Eq.(2) – it leads to an infinite number of solutions of the mass gap equation.

The two cases mostly studied in the literature are the quadratic case,  $\alpha = 2$ , derived in the Gaussian approximation to QCD and using the Balitsky Local Coordinate Gauge [5], and the linear case  $\alpha = 1$ , derived in the Coulomb gauge. In the case of a linear confining potential, we get integral equations. They are finite but need a regularization of infrared divergences.

With the simple harmonic interaction [17], the integral of the potential is a Laplacian, and the mass gap equation can be transformed into a differential equation. The mass gap equation and the quark energy are finally given by,

$$\begin{aligned}\Delta\varphi(k) &= 2kS(k) - 2m_0C(k) - \frac{2S(k)C(k)}{k^2} \\ E(k) &= kC(k) + m_0S(k) - \frac{\varphi'(k)^2}{2} - \frac{C(k)^2}{k^2}.\end{aligned}\quad (5)$$

Numerically, this equation is a non-linear ordinary differential equation. It can be solved with the Runge-Kutta and shooting method.

Examples of solutions, for different light current quark masses  $m_0$ , are depicted in Fig.(5). The effect of a small finite current quark mass  $m_0 \sim 0.01 K_0$ , typical of the light  $u$  and  $d$  quarks, can be easily estimated as a small increase of the dynamically generated constituent quark mass  $m_c$  and does not concern us here.

Examples of replica solutions, for the linear potential, are shown in Fig. 6.

## Introduction

Spontaneous breaking of chiral symmetry and the QCD vacua  
Approach of Gaussian approximation to QCD with two cumulants  
Our previous knowledge of the replicas

## Details of the mass gap and bound state equations

Mass gap equation and replicas

**Salpeter-RPA equations for mesons**

Numerical solution of the mass gap and boundstate equations.

## Summary

	$V^{++} = V^{--}$
spin-indep.	$-\frac{d^2}{dk^2} + \frac{\mathbf{L}^2}{k^2} + \frac{1}{4} (\varphi_q'^2 + \varphi_{\bar{q}}'^2) + \frac{1}{k^2} (\mathcal{G}_q + \mathcal{G}_{\bar{q}})$
spin-spin	$\frac{4}{3k^2} \mathcal{G}_q \mathcal{G}_{\bar{q}} \mathbf{S}_q \cdot \mathbf{S}_{\bar{q}}$
spin-orbit	$\frac{1}{k^2} [(\mathcal{G}_q + \mathcal{G}_{\bar{q}}) (\mathbf{S}_q + \mathbf{S}_{\bar{q}}) + (\mathcal{G}_q - \mathcal{G}_{\bar{q}}) (\mathbf{S}_q - \mathbf{S}_{\bar{q}})] \cdot \mathbf{L}$
tensor	$-\frac{2}{k^2} \mathcal{G}_q \mathcal{G}_{\bar{q}} [(\mathbf{S}_q \cdot \hat{k})(\mathbf{S}_{\bar{q}} \cdot \hat{k}) - \frac{1}{3} \mathbf{S}_q \cdot \mathbf{S}_{\bar{q}}]$
	$V^{+-} = V^{-+}$
spin-indep.	0
spin-spin	$-\frac{4}{3} \left[ \frac{1}{2} \varphi_q' \varphi_{\bar{q}}' + \frac{1}{k^2} \mathcal{C}_q \mathcal{C}_{\bar{q}} \right] \mathbf{S}_q \cdot \mathbf{S}_{\bar{q}}$
spin-orbit	0
tensor	$\left[ -2\varphi_q' \varphi_{\bar{q}}' + \frac{2}{k^2} \mathcal{C}_q \mathcal{C}_{\bar{q}} \right] [(\mathbf{S}_q \cdot \hat{k})(\mathbf{S}_{\bar{q}} \cdot \hat{k}) - \frac{1}{3} \mathbf{S}_q \cdot \mathbf{S}_{\bar{q}}]$

**Table:** The positive and negative energy spin-independent, spin-spin, spin-orbit and tensor potentials are shown.  $\varphi'(k)$ ,  $\mathcal{C}(k)$  and  $\mathcal{G}(k) = 1 - S(k)$  are functions of the quark mass  $m_c(k)$ .

The Salpeter-RPA equations for a meson (a colour singlet quark-antiquark bound state) can be derived either from the  $q\bar{q}$  Lippman-Schwinger equations, or by replacing the propagator of Eq.(3) in the Bethe-Salpeter equation. In either way, one gets [10],

$$\chi(k, P) = \int \frac{d^3 k'}{(2\pi)^3} V(k - k') \left[ u(k'_1) \phi^+(k', P) v^\dagger(k'_2) + v(k'_1) \phi^{-t}(k', P) u^\dagger(k'_2) \right]$$

$$\phi^+(k, P) = \frac{u^\dagger(k_1) \chi(k, P) v(k_2)}{+M(P) - E(k_1) - E(k_2)}, \quad \phi^{-t}(k, P) = \frac{v^\dagger(k_1) \chi(k, P) u(k_2)}{-M(P) - E(k_1) - E(k_2)}. \quad (6)$$

These equations [12, 19] have positive  $\phi^+$  and negative  $\phi^-$  energy wave functions. The Pauli  $\vec{\sigma}$  matrices in the spinors of Eq.(3), produce the [6] potentials of Table 1.

## Introduction


Spontaneous breaking of chiral symmetry and the QCD vacua  
Approach of Gaussian approximation to QCD with two cumulants  
Our previous knowledge of the replicas

## Details of the mass gap and bound state equations


Mass gap equation and replicas  
Salpeter-RPA equations for mesons  
Numerical solution of the mass gap and boundstate equations.

## Summary


Notice that both the pseudoscalar and scalar equations have a system with two equations. However, the spin-dependent interactions couple an extra pair of equations, both in the vector and axial-vector channels. We now combine the spin-dependent potentials of Table 1, to derive the full Salpeter-RPA radial bound state equations.




**Figure:** The constituent quark masses  $m_c(k)$  in the chiral limit  $m_0 = 0$ , solutions of the mass gap equation, from left to right for the ground state vacuum and first two replicas.



**Figure:** The  $J^{PC} = 0^{-+}, {}^1S_0$  normalized pseudoscalar (P) radial wave functions  $\phi^+$  (in blue) and  $\phi^-$  (in yellow), from left to right for the ground state vacuum and first two replicas, in dimensionless units of  $K_0 = 1$ . Because the normalization diverges in the chiral limit, we arbitrarily normalize the wave functions with  $\phi(0) = 1$ . Importantly, the wavefunctions  $\phi^\pm$  are identical to  $\pm \sin \varphi(k) = \pm m_c(k) / \sqrt{k + m_c(k)}$  as is easily verified from Fig. 5.




**Figure:** The  $J^{PC} = 0^{++}, {}^3P_0$  normalized scalar (S) radial wave functions  $\phi^+$  (in blue) and  $\phi^-$  (in yellow), from left to right for the ground state vacuum and first two replicas, in dimensionless in units of  $K_0 = 1$ .




**Figure:** The  $J^{PC} = 1^{--}, {}^3P_1$  normalized vector (V) radial wave functions  $\phi_0^+$  (in blue),  $\phi_0^-$  (in yellow),  $\phi_2^+$  (in green) and  $\phi_2^-$  (in red) from left to right for the ground state vacuum and first two replicas, in dimensionless in units of  $K_0 = 1$ .

To solve numerically our differential equations, we use finite centred differences for the Laplacian. Since our mass gap equation is non-linear, we utilize the shooting method, so as to have the mass to vanish at a large enough UV momentum cutoff  $K$ . We use a very fine mesh in momentum space because the replicas have nodes, quite close to the momentum origin.



**Figure:** The  $J^{PC} = 1^{+-}, ^1P_1$  normalized axial vector (A) radial wave functions  $\phi^+$  (in blue) and  $\phi^-$  (in yellow), from left to right for the ground state vacuum and first two replicas, in dimensionless in units of  $K_0 = 1$ .



**Figure:** The  $J^{PC} = 1^{+-}, ^3P_1$  normalized axial vector (A) radial wave functions  $\phi^+$  (in blue) and  $\phi^-$  (in yellow), from left to right for the ground state vacuum and first two replicas, in dimensionless in units of  $K_0 = 1$ .

Our bound state equations can be understood as an extension of the Schrödinger equation, since we now have positive  $\phi^+$  and negative  $\phi^-$  energy components of the wave functions, and this doubles the number of components. Nevertheless we have the same number of meson states as in the spectrum of the normal quark model. The mass splittings can also be related, as usual, to spin-tensor potentials. We use sparse matrices and matrix eigenvalues to compute the meson spectrum.

We now address the main goal of this study: to show that the quark-antiquark bound states – the mesons in the replicas – all have real excess masses.

With the effective quark masses  $m(k)$ , we can compute the chiral angle, using  $m(k)/k = \tan[\varphi(k)]$ . According to the chiral theorem on the pion Salpeter amplitude [12, 9], the sine of the chiral angle  $\sin[\varphi(k)] = m_c(k)/\sqrt{k + m_c(k)}$  should be proportional to the wave functions  $\phi^\pm(k)$  of the pion. This is clearly the case, when we compare the mass represented in Fig. 5 with the wavefunction shown in Fig. 9 .

For the ground state vacuum and for the first two replicas, in Figs. 9 and 10 , we show, respectively and in dimensionless in units of  $K_0 = 1$ , the wave functions  $\phi^+$  and  $\phi^-$ , for the pseudoscalar meson and the radial wave functions  $\nu^+$  and  $\nu^-$  for the scalar meson. We notice that the number of nodes of these pseudoscalar wave functions depends on the replica where they are sitting, they have the same number of nodes as the constituent quark mass. This is expected due to the chiral theorem relating the pion wave function to the constituent quark mass [12, 9] .

We also show, in Figs. 10 – 13 , the radial wave functions of the scalar, vector and axial vector mesons. Notice how the wave functions change from one replica to another. It is interesting to remark that, for the replicas, the excess masses of the mesons depend very little on the replica (and the physical vacuum) they are sitting in.

The masses scale with  $K_0$ , equivalent to the string tension. Even at finite temperature with a small  $K_0$ , this scenario will hold.

meson	in vacuum	replica $\langle \mathcal{R}1 0 \rangle$	replica $\langle \mathcal{R}2 0 \rangle$	false vacuum
P ( $J^{PC} = 0^{-+}$ )	0.00	0.00	0.00	$i\infty$
P* ( $J^{PC} = 0^{-+}$ )	5.539	5.577	5.581	$i\infty$
S ( $J^{PC} = 0^{++}$ )	3.266	3.253	3.247	$i\infty$
$V_0$ ( $J^{PC} = 1^{--}$ )	2.686	2.823	2.836	3.71
$V_2$ ( $J^{PC} = 1^{--}$ )	4.965	4.635	4.599	4.59
A ( $J^{PC} = 1^{+-}$ )	4.103	3.784	3.723	3.71
A ( $J^{PC} = 1^{++}$ )	4.665	4.602	4.596	4.59

**Table:** Masses of the mesons in the ground state vacuum (the true one) and excess masses for the first two replicas, and false vacuum, in units of  $K_0$ . For the mesons we show the pseudoscalar, the first excitation of the pseudoscalar, the scalar, the vector (mostly s-wave), the first excitation of the vector (mostly d-wave) and the two different axial vectors.

The real nature of the excess masses of the mesons, constitute the main result of this work. Our final results are shown in Table 2.

Prior to this study, we were not sure what the masses of the mesons would be in the excited replica vacua. As in the false - chiral invariant - vacuum, we could possibly have tachyons. Their tachyonic free nature, show that the replicas and thus metastable. More recently, this study has been extended [13] to the linear confining potential, with  $\alpha = 1$  (see Fig.6) and to a small finite current mass  $m_0$  (see Fig. 5) .

As it stands, we cannot have all the low energy properties of hadronic physics due to  $S\chi SB$  without having replicated states as a subproduct. It would be extremely interesting to look whether such metastable replicas do actually exist in full QCD.

### Acknowledgements

P.B. and J.E.R. acknowledge the support of CeFEMA under the FCT contract for R&D Units UIDB/04540/2020.

- [1] A. Amer, A. Le Yaouanc, L. Oliver, O. Pene, and J. c. Raynal. INSTABILITY OF THE CHIRAL INVARIANT VACUUM FOR A CONFINING POTENTIAL. *Phys. Rev. Lett.*, 50:87–90, 1983.
- [2] D. V. Antonov, A. V. Nefediev, and Jose E. F. T. Ribeiro. Dark quark domains. *JETP Lett*, 92:721–725, November 2010.
- [3] P. Bicudo. Mesons and tachyons with confinement and chiral restoration, and NA60. *Phys. Rev.*, D74:065001, 2006.
- [4] P. Bicudo. Quark running mass and vacuum energy density in truncated Coulomb gauge QCD for five orders of magnitude of current masses. *Nucl. Phys.*, A870-871:112–130, 2011.
- [5] P. Bicudo, N. Brambilla, E. Ribeiro, and A. Vairo. Confinement and chiral symmetry breaking in heavy light quark systems. *Phys. Lett.*, B442:349–358, 1998.
- [6] P. J. A. Bicudo, G. Krein, J. E. F. T. Ribeiro, and J. E. Villate. Hadron masses in a chirally symmetric confining model. *Phys. Rev.*, D45:1673–1685, 1992.
- [7] P. J. A. Bicudo and A. V. Nefediev. Chiral symmetry breaking solutions for QCD in the truncated Coulomb gauge. *Phys. Rev.*, D68:065021, 2003.
- [8] P. J. A. Bicudo, J. E. F. T. Ribeiro, and A. V. Nefediev. Vacuum replicas in QCD. *Phys. Rev.*, D65:085026, 2002.
- [9] Pedro Bicudo. Analytic proof that the quark model complies with partially conserved axial current theorems. *Phys. Rev.*, C67:035201, 2003.
- [10] Pedro J. A. Bicudo. A New perspective on the scalar meson puzzle, from spontaneous chiral symmetry breaking beyond BCS. *Phys. Rev.*, C60:035209, 1999.
- [11] Pedro J. de A. Bicudo and Jose E. F. T. Ribeiro. Current Quark Model in a  $p$  Wave Triplet Condensed Vacuum. 1. The Dynamical Breaking of Chiral Symmetry. *Phys. Rev.*, D42:1611–1624, 1990.

- [12] Pedro J. de A. Bicudo and Jose E. F. T. Ribeiro. Current Quark Model in a  $p$  Wave Triplet Condensed Vacuum. 2. Salpeter Equations:  $\pi$ ,  $K$ ,  $\rho$ ,  $\phi$  as  $q\bar{q}$  Bound States. *Phys. Rev.*, D42:1625–1634, 1990.
- [13] Pedro J. de A. Bicudo, Eduardo Garnacho Velasco, Felipe J. Llanes-Estrada, J. Emilio Ribeiro, Víctor Serrano Herreros, and Jorge Vallejo Fernández. Vacuum replicas in field-theory models of Coulomb-gauge QCD. *PoS*, EPS-HEP2021:721, 2022.
- [14] Hans Gunter Dosch and Yu. A. Simonov. The Area Law of the Wilson Loop and Vacuum Field Correlators. *Phys. Lett.*, B205:339–344, 1988.
- [15] L. Ya. Glozman, A. V. Nefediev, and J. E. F. T. Ribeiro. Chiral symmetry restoration in excited hadrons, quantum fluctuations, and quasiclassics. *Phys. Rev.*, D72:094002, 2005.
- [16] Yu. S. Kalashnikova, A. V. Nefediev, and J. E. F. T. Ribeiro. Confinement and parity doubling in heavy-light mesons. *Phys. Rev.*, D72:034020, 2005.
- [17] A. Le Yaouanc, L. Oliver, S. Ono, O. Pene, and J. C. Raynal. A Quark Model of Light Mesons with Dynamically Broken Chiral Symmetry. *Phys. Rev.*, D31:137–159, 1985.
- [18] A. Le Yaouanc, L. Oliver, O. Pene, and J. C. Raynal. Spontaneous Breaking of Chiral Symmetry for Confining Potentials. *Phys. Rev.*, D29:1233–1257, 1984.
- [19] Felipe J. Llanes-Estrada and Stephen R. Cotanch. Meson structure in a relativistic many body approach. *Phys. Rev. Lett.*, 84:1102–1105, 2000.
- [20] Felipe J. Llanes-Estrada and Pedro De A. Bicudo. Quark Schwinger-Dyson evaluation of the L1, L2 coefficients in the chiral Lagrangian. *Phys. Rev.*, D68:094014, 2003.
- [21] Adam P. Szczepaniak and Eric S. Swanson. Coulomb gauge QCD, confinement, and the constituent representation. *Phys. Rev.*, D65:025012, 2002.