



Universitat Autònoma  
de Barcelona



Institut de Física  
d'Altes Energies



# $\eta, \eta' \rightarrow \pi^0 \gamma\gamma$ and $\eta' \rightarrow \eta \gamma\gamma$ decays and a leptophobic U(1) B boson

Rafel Escrivano  
UAB and IFAE-BIST

**QCHS 2022**  
**XVth Quark Confinement and the**  
**Hadron Spectrum Conference**

August 2, 2022  
Stavanger (Norway)



# What's the motivation for this analysis?

## Theoretical analysis of the doubly radiative decays

$$\eta, \eta' \rightarrow \pi^0 \gamma\gamma \text{ and } \eta' \rightarrow \eta \gamma\gamma$$

Decay	Couplings	Chiral loop	$L\sigma M$	VMD	$\Gamma$	$BR_{th}$	$BR_{exp}$ [14]
$\eta \rightarrow \pi^0 \gamma\gamma$ (eV)	Empirical	$1.87 \times 10^{-3}$	$5.0 \times 10^{-4}$	0.16(1)	0.18(1)	$1.35(8) \times 10^{-4}$	$2.56(22) \times 10^{-4}$
	Model-based	$1.87 \times 10^{-3}$	$5.0 \times 10^{-4}$	0.16(1)	0.17(1)	$1.30(1) \times 10^{-4}$	
$\eta' \rightarrow \pi^0 \gamma\gamma$ (keV)	Empirical	$1.1 \times 10^{-4}$	$1.3 \times 10^{-4}$	0.57(3)	0.57(3)	$2.91(21) \times 10^{-3}$	$3.20(7)(23) \times 10^{-3}$
	Model-based	$1.1 \times 10^{-4}$	$1.3 \times 10^{-4}$	0.70(4)	0.70(4)	$3.57(25) \times 10^{-3}$	
$\eta' \rightarrow \eta \gamma\gamma$ (eV)	Empirical	$1.4 \times 10^{-2}$	3.29	21.2(1.2)	23.0(1.2)	$1.17(8) \times 10^{-4}$	$8.25(3.41)(0.72) \times 10^{-5}$
	Model-based	$1.4 \times 10^{-2}$	3.29	19.1(1.0)	20.9(1.0)	$1.07(7) \times 10^{-4}$	

R. Escribano, S. González-Solís, R. Jora and E. Royo, Phys. Rev. D 102 (2020) 034026



# What's a leptophobic U(1) B boson?

It is a **new gauge boson** coupled to the **baryon number**

$$\mathcal{L} = \frac{1}{3} g_B \bar{q} \gamma^\mu q B_\mu$$

$$\alpha_B \equiv g_B^2 / (4\pi)$$

The **low-energy symmetries** of QCD are **preserved**

**C** and **P** are **conserved**

**B** does not transform under **SU(3) flavour symmetry**

**B** is a singlet under **isospin**




$$I^G(J^{PC}) = 0^-(1^{--})$$



**B** is  **$\omega$  meson like!**


# What's a leptophobic U(1) B boson?

B is **not** completely decoupled from leptons



$$\mathcal{L}_{\text{int}} = \left( \frac{1}{3} g_B + \varepsilon Q_q e \right) \bar{q} \gamma^\mu q B_\mu - \varepsilon e \bar{\ell} \gamma^\mu \ell B_\mu$$

with a “natural”-sized  $\varepsilon = eg_B/(4\pi)^2$  induced radiatively



# What's the motivation for a U(1) B boson?

- The **baryon number symmetry** may be related to **dark matter** (it is stabilised since it carries a conserved baryon number charge)
- Natural framework for the **Peccei-Quinn solution** to the **strong CP problem**
- ...

# How are hadronic processes calculated?

Using the **hidden local symmetry (HLS)** for **VMD**

$$\mathcal{L}_{VVP} = \frac{G}{\sqrt{2}} \epsilon^{\mu\nu\alpha\beta} \text{tr} [\partial_\mu V_\nu \partial_\alpha V_\beta P] \quad \text{with} \quad G = \frac{3g^2}{4\pi^2 f_\pi}$$

P is the pseudoscalar meson nonet


V is the vector meson nonet  
**(gauge bosons of a hidden U(3)<sub>V</sub> symmetry)**

In conventional VMD:

$$\mathcal{L}_{V\gamma} = -4egf_\pi^2 A^\mu \text{tr} [QV_\mu]$$

$$\mathcal{L}_{VB} = -4\frac{1}{3}g_B g f_\pi^2 B^\mu \text{tr} [V^\mu]$$

# How are hadronic processes calculated?



# Previous estimates

**Assuming the Narrow Width Approximation (NWA):**

$$\text{BR}(\eta \rightarrow \pi^0 \gamma \gamma) = \text{BR}(\eta \rightarrow B\gamma) \times \text{BR}(B \rightarrow \pi^0 \gamma)$$


and

$$\text{BR}(B \rightarrow \pi^0 \gamma) = 1$$

and **QCD contribution off**

and

$$\text{BR}(\eta \rightarrow \pi^0 \gamma \gamma) < \text{BR}_{\text{exp}} \text{ at } 2\sigma$$



# Present estimates from this analysis


Assuming the **NWA**

and **QCD contribution off**

and

$\text{BR}(\eta \rightarrow \pi^0\gamma\gamma) < \text{BR}_{\text{exp}}$  at  $2\sigma$

and including the  
latest experimental BRs



# However, a lot more can be done nowadays!

Using the new BR value and spectrum from KLOE for

$$\eta \rightarrow \pi^0 \gamma\gamma$$

B. Cao [KLOE-2], PoS EPS-HEP2021 (2022) 409  
E. Pérez del Rio, CD21

Using the recent BR value and spectrum from BESIII for

$$\eta' \rightarrow \pi^0 \gamma\gamma$$

M. Ablikim *et. al.* [BESIII], Phys. Rev. D 96 (2017) 012005

Using the recent BR value from BESIII for

$$\eta' \rightarrow \eta \gamma\gamma$$


M. Ablikim *et. al.* [BESIII], Phys. Rev. D 100 (2019) 052015

# How are these processes calculated?

**VMD:**  $\mathcal{A}_{\eta \rightarrow \pi^0 \gamma \gamma}^{\text{VMD}} = \sum_{V=\rho^0, \omega, \phi} g_{V\eta\gamma} g_{V\pi^0\gamma} \left[ \frac{(P \cdot q_2 - m_\eta^2) \{a\} - \{b\}}{D_V(t)} + \left\{ \begin{array}{l} q_2 \leftrightarrow q_1 \\ t \leftrightarrow u \end{array} \right\} \right]$

**LσM:**  $\mathcal{A}_{\eta \rightarrow \pi^0 \gamma \gamma}^{\text{LσM}} = \frac{2\alpha_{em}}{\pi} \frac{1}{m_{K^+}^2} L(s_K) \{a\} \times \mathcal{A}_{K^+ K^- \rightarrow \pi^0 \eta}^{\text{LσM}}$

**B boson:**  $\mathcal{A}_{\eta \rightarrow \pi^0 \gamma \gamma}^{B \text{ boson}} = g_{B\eta\gamma}(t) g_{B\pi^0\gamma}(t) \left[ \frac{(P \cdot q_2 - m_\eta^2) \{a\} - \{b\}}{D_B(t)} + \left\{ \begin{array}{l} q_2 \leftrightarrow q_1 \\ t \leftrightarrow u \end{array} \right\} \right]$



$$g_{B\pi^0\gamma}(t) = \frac{eg_B}{4\pi^2 f_\pi} F_\omega(t) \quad F_V(s) = \frac{m_V^2}{m_V^2 - s - im_V \Gamma_V}$$


$$g_{B\eta\gamma}(t) = \frac{eg_B}{12\pi^2 f_\pi} \left[ \cos \varphi_P F_\omega(t) + \sqrt{2} \sin \varphi_P F_\phi(t) \right]$$

$$D_B(t) = m_B^2 - t - i\sqrt{t} \Gamma_B(t)$$

# How are these processes calculated?

## B boson width:

$$\begin{aligned}\Gamma_B(t) = & \frac{\tilde{\gamma}_{B \rightarrow e^+ e^-}(t)}{\tilde{\gamma}_{B \rightarrow e^+ e^-}(m_B^2)} \Gamma_{B \rightarrow e^+ e^-} \theta(t - 4m_e^2) + \frac{\tilde{\gamma}_{B \rightarrow \pi^0 \gamma}(t)}{\tilde{\gamma}_{B \rightarrow \pi^0 \gamma}(m_B^2)} \Gamma_{B \rightarrow \pi^0 \gamma} \theta(t - m_{\pi^0}^2) + \frac{\tilde{\gamma}_{B \rightarrow \mu^+ \mu^-}(t)}{\tilde{\gamma}_{B \rightarrow \mu^+ \mu^-}(m_B^2)} \Gamma_{B \rightarrow \mu^+ \mu^-} \theta(t - 4m_\mu^2) \\ & + \frac{\tilde{\gamma}_{B \rightarrow \pi^+ \pi^-}(t)}{\tilde{\gamma}_{B \rightarrow \pi^+ \pi^-}(m_B^2)} \Gamma_{B \rightarrow \pi^+ \pi^-} \theta(t - 4m_\pi^2) + \frac{\tilde{\gamma}_{B \rightarrow \pi^+ \pi^- \pi^0}(t)}{\tilde{\gamma}_{B \rightarrow \pi^+ \pi^- \pi^0}(m_B^2)} \Gamma_{B \rightarrow \pi^+ \pi^- \pi^0} \theta(t - 9m_\pi^2),\end{aligned}$$




# New exclusion plots

Not assuming the NWA

and QCD contribution on

and

$\text{BR}(\eta \rightarrow \pi^0 \gamma\gamma) < \text{BR}_{\text{exp}}$  at  $2\sigma$



$$\text{BR}(\text{PDG}) = (2.56 \pm 0.22) \times 10^{-4}$$


P. A. Zyla et. Al. [PDG], PTEP 2020 (2020) 093C01

$$\text{BR}(\text{KLOE}) = (1.23 \pm 0.14) \times 10^{-4}$$

B. Cao [KLOE-2], PoS EPS-HEP2021 (2022) 409

# New exclusion plots


$$\eta' \rightarrow \pi^0 \gamma\gamma$$



$$\text{BR(BESIII)} = (3.20 \pm 0.07 \pm 0.23) \times 10^{-3}$$

M. Ablikim et. al. [BESIII], Phys. Rev. D 96 (2017) 012005


$$\eta' \rightarrow \eta \gamma\gamma$$




$$\text{BR(BESIII)} = (8.25 \pm 3.41 \pm 0.72) \times 10^{-3}$$

M. Ablikim et. al. [BESIII], Phys. Rev. D 100 (2019) 052015

# Are peaks in the $\pi^0\gamma$ mass distribution seen?




# Fits to the $\gamma\gamma$ mass distribution



**Crystal Ball:**  $\alpha_B = 0.40^{+0.07}_{-0.08}$ ,  $m_B = 583^{+32}_{-20}$  MeV  $\chi^2_{\min}/\text{dof} = 0.4/5 = 0.1$

**KLOE:**  $\alpha_B = 0.049^{+40}_{-27}$ ,  $m_B = 135^{+1}_{-135}$  MeV  $\chi^2_{\min}/\text{dof} = 4.5/5 = 0.9$

# Fits to the $\gamma\gamma$ mass distribution




**BESIII:**  $\alpha_B = 0.005(1)$ ,  $m_B = 759(1)$  MeV  $\alpha_B = 0.018(5)$ ,  $m_B = 156^{+5}_{-1}$  MeV

$$\chi^2_{\text{min}}/\text{dof} = 11.7/11 = 1.1$$

$$\chi^2_{\text{min}}/\text{dof} = 12.5/11 = 1.1$$


# Fits to the $\gamma\gamma$ mass distribution

## Joint Fit



$$\alpha_B = 0.005(1), \quad m_B = 759(1) \text{ MeV}$$

$$\chi^2_{\min}/\text{dof} = 19.6/18 = 1.1$$



$$\alpha_B = 5(2) \times 10^{-4}, \quad m_B = 780^{+3}_{-4} \text{ MeV}$$


$$\chi^2_{\min}/\text{dof} = 23.7/18 = 1.3$$

**BESIII data dominates the fit**


# Conclusions

- The sensitivity of the rare decays  $\eta, \eta' \rightarrow \pi^0\gamma\gamma$  and  $\eta' \rightarrow \eta\gamma\gamma$  to a leptophobic U(1) B boson in the mass range MeV-GeV has been analysed in detail
- Stringent limits on the B boson parameters  $m_B$  and  $a_B$  have been found
- The current constraints have been strengthened by one order of magnitude from  $\eta \rightarrow \pi^0\gamma\gamma$
- These constraints would make a B-boson signature strongly suppressed





# Conclusions



# Backup slides



# Backup slides



# Backup slides

