Massive secondary quark corrections to bHQET cross section

Alejandro Bris¹

in colaboration with Vicent Mateu, Fernando Gil

¹Universidad Autónoma de Madrid

XVth Quark confinement and the Hadron spectrum Stavanger, August 2 2022

Outline

Motivation

2 Massive secondary quark bHQET

- Massless Factorization Theorem
- Mass corrections

Matrix Elements

- Dispersive Integral Method
- Jet Function
- bHQET Hard Function

Flavour Matchings

- Jet Function
- bHQET Hard Function

Motivation

I Precise determination of top quark mass needs to account for bottom mass efects:

Observable sensible to the quark mass. Hemisphere Jet mass:

$$s \equiv \left(\sum_{i \in h} p_i\right)^2$$
 $s_{\min} = M_t^2$

Extract the primary quark mass from the peak position of the distribution



- $\diamond~$ Close to the peak, fluctuations around the primary quark mass are very small: $\frac{s-M_t^2}{M_t} \ll M_t$
- $\diamond~$ A non-vanishing secondary quark mass is relevant for a precise determination of the peak position
- II Massive secondary quark corrections might play important role for bottom mass determinations as well
- III These corrections are the missing piece of $\mathcal{O}(\alpha_s^2)$ computations

Dijets' primary quarks production



* Q center-of-mass energy



- 1) Integrate out the heavy quark and antiquark masses in their corresponding rest frames \Rightarrow $2\times$ HQET's
- 2) Boost back to c.o.m. frame
- 3) Match onto SCET in order to account for global soft radiation.

	REST FRAMES		BOOSTED FRAME (bHQET)	
	DOF	Scalings	DOF	Scalings
HQET ₁	quark: $p = Mv + k$	v = (1, 1, 0)	quark: $p = Mv_+ + k_+$	$v_+ = \left(rac{M}{Q}, rac{Q}{M}, 0 ight)$
	soft: k	$k \sim \Gamma(1, 1, 1)$	<i>n</i> -ucollinear: k_+	$k_+ \sim \Gamma\left(rac{M}{Q}, rac{Q}{M}, 1 ight)$
HQET ₂	anti-quark: $p = Mv + k$	v=(1,1,0)	anti-quark: $p = Mv + k$	$v_{-} = \left(rac{Q}{M}, rac{M}{Q}, 0 ight)$
	soft: <i>k</i>	$k \sim \Gamma(1, 1, 1)$	\bar{n} -ucollinear: k_{-}	$k_{-} \sim \Gamma\left(rac{Q}{M},rac{M}{Q},1 ight)$
			soft: q _s	$q_{s} \sim rac{\Gamma M}{Q}\left(1,1,1 ight)$

* M primary quark mass

bHQET factorization theorem (e^+e^- production)

Integrate out hard momentum top-quark shells at scale μ_M

* 2-jettiness (Thrust M-scheme):

$$\begin{split} \tau \equiv & \frac{1}{Q} \min_{\hat{t}} \sum_{i} (p_{i}^{0} - |\hat{t} \cdot \vec{p}_{i}|) \\ \tau \approx & \frac{s_{h^{+}} + s_{h^{-}}}{Q^{2}} \quad \text{in the peak region} \end{split}$$



$$\frac{1}{\sigma_0} \frac{\mathsf{d}\hat{\sigma}_{\mathrm{bHQET}}}{\mathsf{d}\tau} = Q^2 H(Q, \mu_M) H_M\left(M, \frac{Q}{M}, \mu_M, \mu\right) \int \mathsf{d}\ell B_{\tau}\left(\frac{Q^2(\tau - \tau_{\min}) - Q\ell}{M}, \mu\right) S_{\tau}(\ell, \mu)$$

$$B_{\tau}(\hat{s},\mu) = M \int_0^{\hat{s}} \mathrm{d}\hat{s}' B_n(\hat{s}-\hat{s}',\mu) B_n(\hat{s}',\mu)$$

- $*~~{\it H}({\it Q},\mu_{\it M})$ and ${\it S}_{ au}(\ell,\mu)$ same as SCET
- * $S_{\tau}(\ell,\mu)$ massive secondary quark corrections already known: [S. Gritschacher, A.Hoang, I.Jemos, P. Pietrulewicz, 2013]

bHQET

Massive secondary quark corrections

New scale (secondary quark mass) brings a richer structure of EFTs:

 \rightarrow Different scenarios:



b) is the relevant in the peak region

ightarrow Consistency conditions: freedom to choose scale at which everyone runs to.



Dispersive Integral Method

Method:

- 1) Write massive bubble diagram as an integral of an effective gluon propagator
- 2) Perform computations at previous loop order with the modified gluon propagator
- 3) Carry out dispersive integral

MASSIVE GLUON:

$$\begin{aligned} \text{MASSIVE BUBBLE} &\longrightarrow \frac{-i}{p^2} \left(g^{\mu\nu} - \frac{p^{\mu}p^{\nu}}{p^2} \right) \Pi(p^2, m^2) \\ \frac{\Pi(p^2, m^2) - \Pi(0, m^2)}{p^2} &= T_F \frac{\alpha_s}{4\pi} \int_{4m^2}^{\infty} \mathrm{d}\tilde{m}^2 \frac{\mathcal{V}^{(d)}(\tilde{m}, \mu)}{p^2 - \tilde{m}^2 + i\epsilon} \\ \mathcal{V}^{(d)}(\tilde{m}, \mu) &= \frac{8\Gamma(2-\varepsilon)}{\Gamma(4-2\varepsilon)} \frac{\beta_{\tilde{m}}}{\tilde{m}^2} \left(\frac{4\pi\tilde{\mu}^2}{\beta_{\tilde{m}}^2 \tilde{m}^2} \right)^{\varepsilon} \left(1 - \varepsilon + \frac{2M^2}{\tilde{m}^2} \right) \end{aligned}$$

 m bubble quark mass, $^\tilde{m}$ gluon effective mass, $^*\beta_{\tilde{m}}\equiv\sqrt{1-4M^2/\tilde{m}^2}$

Using Mellin-Barnes representation for the modified gluon propagator reduces complexity to massless 1-loop with modified exponents. Same computation as in renormalon calculus

- MELLIN PLANE:
 - Closed expression for $\Pi(p^2, m^2)$ obtained from massive gluon integration:

$$\Pi(p^2,m^2) = \frac{T_F \,\Gamma(\varepsilon)}{2\varepsilon - 3} \frac{\alpha_s}{\pi} \left(\frac{4\pi\tilde{\mu}^2}{m^2}\right)^{\varepsilon} \left[\left(1 - \varepsilon + \frac{2m^2}{p^2}\right) \,_2F_1\left(1,\varepsilon;\frac{3}{2};\frac{p^2}{4m^2}\right) - \frac{2m^2}{p^2} \right]$$

II Integral representation of $_2F_1$:

$${}_{2}F_{1}\left(1,\varepsilon;\frac{3}{2};\frac{p^{2}}{4m^{2}}\right) = \frac{\Gamma(3/2)}{\Gamma(\varepsilon)\,\Gamma(3/2-\varepsilon)}\int_{0}^{1}\mathrm{d}x\frac{x^{-1+\varepsilon}(1-x)^{1/2-\varepsilon}}{1-\frac{p^{2}}{4m^{2}}x}$$

III Mellin-Barnes representation:

$$\frac{1}{1-\frac{p^2}{4m^2}x} = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} \mathrm{d}h \, \left(-\frac{p^2}{4m^2}x\right)^{-h} \Gamma(h) \, \Gamma(1-h)$$

IV Integration of x variable can be carried out right away

After integrating \times one finds:

Mellin Plane Result

$$\frac{\Pi(p^2,m^2)-\Pi(0,m^2)}{p^2} = \frac{T_F \alpha_s}{2\pi^2 i p^2} \left(\frac{4\pi \tilde{\mu}^2}{m^2}\right)^{\varepsilon} \int_{c-i\infty}^{c+i\infty} \mathrm{d}h \left(-\frac{m^2}{p^2}\right)^{-h} \frac{1+h}{3+2h} \frac{h\Gamma^2(h)\Gamma(1-h)\Gamma(h+\varepsilon)}{\Gamma(2h+2)}$$

 \Rightarrow Modified gluon propagator: same as renormalon calculus "analytic regulator"

$$rac{-ig^{\mu
u}}{p^2} \longrightarrow rac{ig^{\mu
u}}{(-p^2)^{1-h}}$$

- st Does not introduce an additional energy scale in previous order computations \checkmark
- * Integration of h by residues yields directly expansions for large or small mass (converse mapping theorem) \checkmark
- ∗ It also may give rise to closed forms in terms of Meijer functions ✓
- Dealing with distributions needs extra effort X

Definition:

$$\mathcal{B}_{n}(\hat{s},\mu) = \frac{i}{4\pi N_{C}M} \operatorname{Tr}\left[\int d^{d}x \, e^{ikx} \left\langle 0 \right| T\left\{ W_{n}^{+}(x) \, h_{v}(x) \, \bar{h}_{v}(0) \, W_{n}(0) \right\} \left| 0 \right\rangle \right]$$
$$B_{n}(\hat{s},\mu) = \operatorname{Im}\left[\mathcal{B}_{n}(\hat{s},\mu)\right]$$

- * W_n Wilson lines with u-collinear gluons
- * h_v Heavy quark field

Contributing diagrams:



One loop results. Massive gluon

Calculation comments:

- $\rightarrow\,$ Rapidity divergence appears in vertex diagram due to non-zero gluon mass \Rightarrow Introduce $\delta\,$ regulator that changes de exponent.
- $\rightarrow \mathcal{B}_n(\hat{s}, \mu, \tilde{m})$ involves hypergeometric functions with \hat{s} dependence which will lead to "hidden" distributions when taking the imaginary part \Rightarrow Easiest way:

$$\mathsf{Im}\Big[\mathcal{B}_n(\hat{s},\mu,\tilde{m})\Big] = \mathsf{Im}\Big[\mathcal{B}_n(\hat{s},\mu,\tilde{m}\to\infty)\Big] + \mathsf{Im}\Big[\mathcal{B}_n(\hat{s},\mu,\tilde{m}) - \mathcal{B}_n(\hat{s},\mu,\tilde{m}\to\infty)\Big]_{\varepsilon\to0} + \mathcal{O}(\varepsilon)$$

ightarrow regulator dependence disappears when taking imaginary part

$B_n(\hat{s},\mu,\tilde{m})$

$$\begin{split} &\frac{\alpha_{s}C_{F}}{4\pi M} \left\{ 2\,\Gamma(\varepsilon) \left(\frac{\mu^{2}}{\tilde{m}^{2}}e^{\gamma}\right)^{\varepsilon} \Bigg[\left(-H_{\varepsilon-1}+2\log(\tilde{m})+1\right) \delta(\hat{s}) - 2\left[\frac{\theta(\hat{s})}{\hat{s}}\right]_{+} - \frac{2\tilde{m}\pi^{1/2}\,\Gamma(1/2+\varepsilon)}{(2\varepsilon-1)\,\Gamma(\varepsilon)}\,\delta'(\hat{s}) \Bigg] \right. \\ &\left. + \theta\left(\hat{s}^{2}-4\tilde{m}^{2}\right) \left[\frac{8}{\hat{s}}\log\left(\frac{\hat{s}+\sqrt{\hat{s}^{2}-4\tilde{m}^{2}}}{2\tilde{m}}\right) - \frac{4\sqrt{\hat{s}^{2}-4\tilde{m}^{2}}}{\hat{s}^{2}} \right] + \mathcal{O}(\varepsilon) \right\} \end{split}$$

reproduces massless result

Calculation comments:

- ightarrow Simpler integrals than massive gluon case because there is only one scale: \hat{s}
- \rightarrow Rapidity regulator is not needed
- \rightarrow Imaginary part of just powers:

$$(-\hat{s}-i\epsilon)^{-a}=\hat{s}^{-a}e^{ai\pi}\xrightarrow{i\mathrm{Im}}\hat{s}^{-a}\frac{i\pi}{\Gamma(a)\Gamma(1-a)},$$

$B_n^h(\hat{s},\mu)~(\equiv$ One loop jet function with gluon propagator $rac{ig^{\mu u}}{(-p^2)^{1-h}}$)

$$-C_{\mathsf{F}}\frac{\alpha_{\mathsf{s}}}{\pi M}\frac{\Gamma(2+h-\varepsilon)\,\hat{\mathsf{s}}^{-1+2h}e^{\varepsilon\gamma}}{(\varepsilon-h)\Gamma(1-h)\Gamma(2+2h-2\varepsilon)}\left(\frac{\mu}{\hat{\mathsf{s}}}\right)^{2\varepsilon}$$

no need to expand in epsilon or h

agreement with [N. G. Gracia, V. Mateu, 2021]

Two loop massive secondary quark corrections. Massive gluon Computation

All terms in dispersive integral analytically computed easily except:

$$I[\hat{s}, m] \equiv \frac{1}{4\hat{s}} \int_{4m^2}^{\hat{s}^2/4} \mathrm{d}\tilde{m}^2 \, \, \mathcal{V}^{(4)}\left(\tilde{m}, \mu\right) \, \log\left(\frac{\hat{s} + \sqrt{\hat{s}^2 - 4\tilde{m}^2}}{2\tilde{m}}\right)$$

Integrating by parts and defining $a \equiv \sqrt{1 - \frac{16m^2}{s^2}}$ we managed to obtain an expression in terms of hypergeometric functions and elliptic integrals *E* and *K*:

$$I[\hat{s}, m] = \frac{1}{3\hat{s}} \left[\frac{1}{9} \left(16 - 2\hat{s}^2 \right) E\left(\hat{s}^2 \right) - \frac{1}{9} \left(16 - \hat{s}^2 \right) K\left(\hat{s}^2 \right) + f(\hat{s}) \right]$$

$$\begin{split} f(a) &= -\frac{1}{8} \left(1-a^2\right) \, {}_5F_4\left(1,1,1,\frac{3}{2},\frac{3}{2};2,2,2,2;1-a^2\right) \\ &+ \left[\frac{1-a^2}{4}+\frac{1-a^2}{8} \log\left(\frac{1-a^2}{16}\right)\right] \, {}_4F_3\left(1,1,\frac{3}{2},\frac{3}{2};2,2,2;1-a^2\right) \\ &+ \frac{1-a^2}{4} \frac{d}{d\epsilon} \left[\, {}_4F_3\left(1,1,\frac{3}{2},\frac{3}{2}+\epsilon;2,2,2+\epsilon;1-a^2\right) \right]_{\epsilon=0} + \frac{1}{4} \log^2\left(\frac{1-a^2}{16}\right) - \frac{\pi^2}{6} \end{split}$$

Two loop massive secondary quark corrections. Mellin plane Computation

$$\begin{split} B_n^{m-\text{bubble},OS}\left(\hat{s},\mu,m\right) &= \frac{T_f \, \alpha_s}{2\pi^2 i \, M} \left(\frac{4\pi \tilde{\mu}^2}{m^2}\right)^{\varepsilon} \int_{c-i\infty}^{c+i\infty} dh \left(m^2\right)^{-h} \frac{1+h}{3+2h} \frac{h \, \Gamma^2(h)\Gamma(1-h)\Gamma(h+\varepsilon)}{\Gamma(2h+2)} \, B_n^h\left(\hat{s},\mu\right) \\ &= -\frac{\alpha_s^2 \, T_f \, C_F}{2^{4-2\varepsilon} \, \pi^2 i \, M} \left(\frac{\mu^2 e^{\gamma}}{m}\right)^{2\varepsilon} \hat{s}^{-1-2\varepsilon} \int_{c-i\infty}^{c+i\infty} dh \left(\frac{16m^2}{\hat{s}^2}\right)^{-h} \frac{(1-\varepsilon+h)(1+h) \, \Gamma(h)\Gamma(h+\varepsilon)}{(\varepsilon-h) \, \Gamma(\frac{5}{2}+h)\Gamma(\frac{3}{2}-\varepsilon+h)} \\ &= -\frac{\alpha_s^2 \, T_f \, C_F}{2^{3-2\varepsilon} \, \pi \, M} \left(\frac{\mu^2 e^{\gamma}}{m}\right)^{2\varepsilon} \hat{s}^{-1-2\varepsilon} \, G_{5,5}^{4,1} \left(\frac{16m^2}{\hat{s}^2}\right| \frac{1-\varepsilon, 1, 1-\varepsilon, \frac{5}{2}, \frac{3}{2}-\varepsilon}{0, \varepsilon, 2, 2-\varepsilon, -\varepsilon} \end{split}$$

* Trick: one can postpone MB inversion after resummation

• Converse mapping theorem:

Fundamental strip:
$$h \in (0, \varepsilon)$$
. Expansion series: $\begin{cases} \text{l.h.s poles: } rac{16m^2}{s^2} \to 0 \\ \text{r.h.s poles: } rac{16m^2}{s^2} \to \infty \end{cases}$

Distributional structure:

$$B_{n}^{m-\text{bubble},OS}\left(\hat{s},\mu,m\right) \equiv f_{\delta} \ \delta(\hat{s}) + \sum_{i=0} f_{\mathcal{L}_{i}} \left[\frac{\theta(\hat{s})\log^{i}(\hat{s})}{\hat{s}}\right]_{+} + \sum_{j=1} f_{\delta(j)} \ \delta^{(j)}(\hat{s}) + \theta\left(g(\hat{s},m)\right) f_{nd}(\hat{s}) + \theta\left(g(\hat{s},m)\right) \left(g(\hat{s},m)\right) \left(g(\hat{s},m)\right) + \theta\left(g(\hat{s},m)\right) \left(g(\hat{s},m)\right) \left(g(\hat{s},m)\right) + \theta\left(g(\hat{s},m)\right) \left(g(\hat{s},m)\right) \left(g(\hat{s},m)\right) + \theta\left(g(\hat{s},m)\right) \left(g(\hat{s},m)\right) \left(g(\hat{s}$$

$$\delta^{(j)}(x) \equiv \frac{\mathrm{d}^j}{\mathrm{d}x^j} \delta(x)$$

Final Result

Carrying out renormalization and convolving both hemisphere jet functions we get:

$$\begin{split} \mathfrak{g}_{\tau}^{(n_{l}+1)}(\hat{s},\mu,m) &= \mathcal{B}_{\tau}^{(n_{l}+1)}(\hat{s},\mu) + \delta \mathcal{B}_{m}^{dist}(\hat{s},\mu,m) + \delta \mathcal{B}_{m}^{real}(\hat{s},m) \\ \delta \mathcal{B}_{m}^{dist}(\hat{s},\mu,m) &= \left(\frac{\alpha_{s}^{(n_{l}+1)}}{4\pi}\right)^{2} \frac{C_{F}}{M} \left[\left(\frac{32}{9}L_{m}^{3} + \frac{128}{9}L_{m}^{2} + \left(\frac{976}{27} - \frac{16\pi^{2}}{9}\right)L_{m} + \frac{3568}{81} - \frac{64\pi^{2}}{27} - \frac{32}{3}\xi_{3} \right) \delta(\hat{s}) \\ &+ \left(-\frac{32}{3}L_{m}^{2} - \frac{256}{9}L_{m} - \frac{976}{27} + \frac{16\pi^{2}}{9} \right) \mathcal{L}^{0}(\hat{s}) + \left(\frac{64}{3}L_{m} + \frac{256}{9}\right)\mathcal{L}^{1}(\hat{s}) - \frac{32}{3}\mathcal{L}^{2}(\hat{s}) - 8\pi^{2}m\,\delta'(\hat{s}) \right] \\ \delta \mathcal{B}_{m}^{real}(\hat{s},m) &= \left(\frac{\alpha_{s}^{(n_{l}+1)}}{4\pi}\right)^{2}\frac{C_{F}}{M}\frac{\theta\left(\hat{s}^{2} - 16m^{2}\right)}{\hat{s}} \left[\frac{976}{27} - \frac{16\pi^{2}}{9} + \frac{256}{9}\log\left(\frac{m}{\hat{s}}\right) + \frac{32}{3}\log^{2}\left(\frac{m}{\hat{s}}\right) \\ &+ \sum_{n=0}^{\infty}\frac{\left(\left(\frac{1}{2}\right)_{n}\right)^{2}}{(n!)^{2}(n+2)^{3}}\left(\frac{4n-1}{n+2} + 2(1+2n)\left[\psi(n+1) - \psi(n+1/2) - \log\left(\frac{4m}{\hat{s}}\right)\right]\right)\left(\frac{16m^{2}}{\hat{s}^{2}}\right)^{2+n} \right] \end{split}$$

 $-B_{ au}^{(n_l+1)}(\hat{s},\mu)$ can be found in [A. Jain, I. Scimemi and I.W. Stewart, 2008]

- n_l number of massless flavours
- $\begin{array}{l} \ L_m \equiv \log\left(\frac{m}{\mu}\right) \\ \\ \ \mathcal{L}^i(\hat{s}) \equiv \frac{1}{\mu} \left[\frac{\theta(\hat{s})\log^i(\hat{s}/\mu)}{\hat{s}/\mu}\right]_+ \end{array}$

One loop results

Definition:

$$\mathcal{J}_{\mathsf{SCET}} = \mathcal{C}_M \, \mathcal{J}_{\mathsf{bHQET}}$$

 $\mathcal{H}_M = |\mathcal{C}_M|^2$

Computation:

$$C_{M} = \frac{\langle q, \bar{q} | \mathcal{J}_{\mathsf{SCET}} | 0 \rangle}{\langle q, \bar{q} | \mathcal{J}_{\mathsf{bHQET}} | 0 \rangle} = \frac{F_{\mathsf{SCET}}}{F_{\mathsf{bHQET}}}$$

Contributing diagrams:



One loop results

Massive gluon

 $F_{SCET}^{\tilde{m}} \rightarrow$ [A. H. Hoang, A. Pathak, P. Pietrulewicz, I. W. Stewart, 2015]

$$F_{\rm bHQET}^{\tilde{m}} = \frac{\alpha_{\rm s} C_{\rm F}}{2\pi} \Gamma(\varepsilon) \left(\frac{\mu^2}{\tilde{m}^2} {\rm e}^{\gamma}\right)^{\varepsilon} \left[1 + \log\left(\frac{M^2}{Q^2}\right) + i\pi\right]$$

• Mellin plane

$$F_{\text{SCET}}^{h} = \frac{\alpha_{s}C_{F}}{\pi} \left(\mu^{2}e^{\gamma}\right)^{\varepsilon} \left(h-1\right) \left(1 + \frac{(3-2\varepsilon)(\varepsilon-h-1)(\varepsilon-h)}{2+h-2\varepsilon}\right) \frac{\Gamma(\varepsilon-h)\Gamma(2h-2\varepsilon)}{\Gamma(2+h-2\varepsilon)} \left(M^{2}\right)^{h-\varepsilon}$$
$$F_{\text{bHQET}}^{h} \text{ Scaleless} \to C_{M}^{h} = F_{\text{SCET}}^{h}$$

Final result for expansion

$$C_{M}^{(n_{l}+1)}\left(M,\frac{Q}{M},\mu,m\right) = C_{M}^{(n_{l}+1)}\left(M,\frac{Q}{M},\mu\right) + \delta C_{M}\left(\frac{m}{M}\right)$$

$$\begin{split} \delta C_M \left(\frac{m}{M}\right) &= \left(\frac{\alpha_s^{(n_f+1)}}{4\pi}\right)^2 C_F T_f \left[6\pi^2 \frac{m}{M} + \frac{4m^2}{M^2} \left(6 + 8\hat{L}_m \right) - \frac{110}{9} \pi^2 \frac{m^3}{M^3} \right. \\ &+ \frac{m^4}{3M^4} \left(145 + 12\pi^2 - 72 \left(2 - \hat{L}_m \right) \hat{L}_m \right) + \frac{m^6}{M^6} \sum_{n=0}^\infty a_n (m/M) \left(\frac{m^2}{M^2} \right)^n \right] \end{split}$$

$$\begin{aligned} a_n(m/M) &= \frac{8}{(n+1)(n+2)(n+3)^3(2n+3)(2n+5)} \Biggl[506 + 750n + 413n^2 + 100n^3 + 9n^4 \\ &+ \left(408 + 778n + 589n^2 + 221n^3 + 41n^4 + 3n^5 \right) \log(4) + (n+2)(n+3)(n+4) \left(17 + 14n + 3n^2 \right) \times \\ &\times \Biggl[2 \, \hat{L}_m + \psi(-1/2 - n) + \psi(1+n) - 2 \, \psi(7+2n) \Biggr] \Biggr] \end{aligned}$$

- $-C_M^{(n_l+1)}\left(M, \frac{Q}{M}, \mu\right)$ can be found in [A. H. Hoang, A. Pathak, P. Pietrulewicz, I. W. Stewart, 2015]
- n_l number of massless flavours
- $-\hat{L}_m \equiv \log\left(\frac{m}{M}\right)$

Final result unexpanded

$$\begin{split} \delta C_M\left(\frac{m}{M}\right) &= \left(\frac{\alpha_s^{(n_f+1)}}{4\pi}\right)^2 C_F T_f \left\{\frac{1747}{81} + \frac{52\pi^2}{27} + \left(\frac{532}{27} + \frac{16}{9}\pi^2\right)\hat{L}_m + \frac{104}{9}\hat{L}_m^2 + \frac{32}{9}\hat{L}_m^3 \right. \\ &+ \frac{1}{2}\int_{4m^2}^{\infty} \mathrm{d}\hat{m}^2 \,\mathcal{V}^{(4)}\left(\tilde{m},\mu\right) \left[\frac{\left(-b^3 - 5b + 6\right)}{b(b+1)^2}\log\left(\frac{1-b}{2}\right) - \frac{\left(b^3 + 5b + 6\right)}{(1-b)^2b}\log\left(\frac{b+1}{2}\right) \right. \\ &+ \frac{b^2 - 25}{2\left(1-b^2\right)} + 4\log\left(\frac{1-b}{2}\right)\log\left(\frac{b+1}{2}\right) \right] \right\} \end{split}$$

$$- b \equiv \sqrt{1 - \frac{4M^2}{\tilde{m}^2}}$$

n_l number of massless flavours

$$-\hat{L}_m \equiv \log\left(\frac{m}{M}\right)$$

Based on [P. Pietrulewicz, S. Gritschacher, A. H. Hoang, I. Jemos, V. Mateu, 2014]

$$\mathcal{M}_B = M \int \mathrm{d}\hat{s}' \; B^{(n_l)}_{ au}(\hat{s} - \hat{s}', \mu_m, m) \left[B^{(n_l+1)}_{ au}(\hat{s}', \mu_m, m)
ight]^{-1}$$

$$B_{ au}^{(n_l)}(\hat{s},\mu,m) = B_{ au}^{(n_l)}(\hat{s},\mu) + B_{ au}^{m- ext{bubble,OS}}\left(\hat{s},\mu,m
ight) - Z_B^{OS}\left(\hat{s},\mu,m
ight)$$

 $Z_B^{OS}(\hat{s}, \mu, m)$ is obtained from the decoupling condition:

$$\lim_{m\to\infty} \left[B_{\tau}^{(n_l)}(\hat{s},\mu,m) \right] = B_{\tau}^{(n_l)}(\hat{s},\mu) \underbrace{- \left(\frac{\alpha_s}{4\pi}\right)^2 \frac{C_F}{M} 8\pi^2 m \, \delta'(\hat{s})}_{\diamondsuit}$$

> comes from the Lagrangian of bHQET^(n_l) theory

Flavour Matching Jet Function

$$\mathcal{L}_{bHQET}^{(n_{l}+1)} = \bar{h}_{v} (i v \cdot D - \delta M) h_{v}$$

$$\delta M = M_{pole} - M$$

$$\bigcup$$

$$\mathcal{L}_{bHQET}^{(n_{l})} = \bar{h}_{v} \left[i v \cdot D - \delta M - \left(\frac{\alpha_{s}}{4\pi}\right)^{2} C_{F} 2\pi^{2} m \right] h_{v}$$

• Massive secondary quark bubbles remain as a contribution to the primary quark self-energy

$$\mathcal{M}_{B}(\hat{s},\mu_{m},m) = \left(\frac{\alpha_{s}^{(n_{l}+1)}}{4\pi}\right)^{2} \frac{C_{F}}{M} \left\{ \left[-\frac{32}{9}L_{m}^{3} - \frac{128}{9}L_{m}^{2} + \left(-\frac{688}{27} + \frac{4\pi^{2}}{9}\right)L_{m} - \frac{440}{27} + \frac{5\pi^{2}}{27} + \frac{28}{9}\xi_{3} \right] \delta(\hat{s}) + \left(\frac{32}{3}L_{m}^{2} + \frac{160}{9}L_{m} + \frac{224}{27}\right)\mathcal{L}^{0}(\hat{s}) \right\}$$

Flavour Matching bHQET Hard Function

$$\mathcal{M}_{C_{M}} = \frac{C_{M}^{(n_{l})}\left(M, \frac{Q}{M}, \mu_{m}, m\right)}{C_{M}^{(n_{l}+1)}\left(M, \frac{Q}{M}, \mu_{m}, m\right)} \quad ; \qquad \qquad \mathcal{M}_{H_{M}} = \left|\mathcal{M}_{C_{M}}\right|^{2}$$

$$C_{M}^{(n_{l})}\left(M,\frac{Q}{M},\mu,m\right) = C_{M}^{(n_{l})}\left(M,\frac{Q}{M},\mu\right) + C_{M}^{m-\text{bubble},OS}\left(M,\mu,m\right) - Z_{C_{M}}^{OS}\left(M,\mu,m\right)$$

*~ Decoupling limit \Rightarrow Remove the quark also from the running of SCET hard function:

$$\lim_{m \to \infty} \left[H^{(n_l+2)}(Q,\mu,m) H_M^{(n_l)} \left(M, \frac{Q}{M}, \mu, m \right) \right] = H^{(n_l+1)}(Q,\mu) H_M^{(n_l)} \left(M, \frac{Q}{M}, \mu \right)$$

Flavour Matching bHQET Hard Function

2

$$H^{(n_l+1)}(Q,\mu) = \lim_{m \to \infty} \left[|\mathcal{M}_C(Q,\mu,m)|^2 \ H^{(n_l+2)}(Q,\mu,m) \right]$$
$$\bigcup_{\text{Re} \left[C_M^{m-\text{bubble},OS}(M,\mu,m) - Z_{C_M}^{(2),OS}(M,\mu,m) - \mathcal{M}_C^{(2)}(Q,\mu,m) \right]} \xrightarrow[m \to \infty]{} 0$$

 $\mathcal{M}_{\mathcal{C}}(\mathcal{Q},\mu,m) \rightarrow [P. Pietrulewicz, S. Gritschacher, A. H. Hoang, I. Jemos, V. Mateu, 2014]$

$$\mathcal{M}_{H_M}^{(2)}\left(M,\frac{Q}{M},\mu_m,m\right) = \left(\frac{\alpha_s^{(n_l+1)}}{4\pi}\right)^2 C_F \frac{16}{27} \left[9\log^2\left(\frac{m}{\mu_m}\right) + 15\log\left(\frac{m}{\mu_m}\right) + 7\right] \left[1 - 2\log\left(\frac{Q}{M}\right)\right]$$

Numerical Results

Based on [B. Bachu, A. H. Hoang, V. Mateu, A. Pathak, I. W. Stewart, 2020]

 $M_t^{
m pole} = 170.034$ Q = 2000 $\Gamma_t = 1.32$



 $\rightarrow\,$ For a precise determination of the peak position, massive secondary quark corrections need to be taken into account.

 $\rightarrow\,$ Missing pieces of thrust and hemisphere mass distribution in the bHQET factorization theorem were computed.

 $\rightarrow\,$ We proposed a dispersive integration method in Mellin plane to compute massive bubble corrections.

 \rightarrow The results were implemented in a numerical code for thrust bHQET cross section at N³LL + $O(\alpha_s^2)$ accuracy.

BACKUP SLIDES

Two loop massive secondary quark corrections. Mellin plane Computation

Recovering distributions:

* I.h.s poles
$$\Rightarrow \sum_{i=0} f_{\mathcal{L}_i} \frac{\log^i(\hat{s})}{\hat{s}} + f_{nd}$$

- * r.h.s pole (ONLY $h = \varepsilon$) $\Rightarrow \sum_{i=0} f_{\mathcal{L}_i} \frac{\log^i(\hat{s})}{\hat{s}}$
- * Dirac delta and its derivatives:

$$\int_{0}^{\hat{s}_{c}} d\hat{s} \, \hat{s}^{k} B_{n}^{m-\text{bubble},OS}\left(\hat{s},\mu,m\right) \bigg|_{\mathcal{O}\left(\hat{s}_{c}^{0}\right)} = (-1)^{k} k! f_{\delta^{(k)}}$$

Different ways:

- * use $\hat{s}^{-1+2(h-\varepsilon)} = \frac{1}{2(h-\varepsilon)} \,\delta(\hat{s}) + \sum_{n=0} \frac{2^n (h-\varepsilon)^n}{n!} \left[\frac{\theta(\hat{s}) \log^n(\hat{s})}{\hat{s}}\right]_+$ for dirac delta and plus distributions
- * Setting $\varepsilon = 0$ directly in the Mellin plane expression one recovers the non-distributional part as a Meijer G-function:

$$\theta \left(1 - \frac{16m^2}{\hat{s}^2} \right) f_{\mathsf{nd}} = \frac{\alpha_s^2 \, T_f \, C_F}{8\pi \, M} \hat{s}^{-1} \, G_{5,5}^{5,0} \left(\frac{16m^2}{\hat{s}^2} | \begin{array}{c} 1, 1, 1, \frac{3}{2}, \frac{5}{2} \\ 0, 0, 0, 2, 2 \end{array} \right)$$

* Obtain the Mellin plane two loops expression for $\mathcal{B}_n^{m-\text{bubble},OS}(\hat{s},\mu,m)$, expand in $\hat{s} = \hat{s} + i\epsilon$ and take then the imaginary part