

*New birational invariants

Ludmil Katzarkov - University of Miami

Birational Geometry

An algebraic variety X is *rational* over \mathbb{C} if $\mathbb{C}(X) \cong \mathbb{C}(x_1, \dots, x_n)$, where $n = \dim_{\mathbb{C}} X$.

Example

Let $X \subset \mathbb{P}^2$ be a smooth cubic curve.

- * Hodge numbers: $h^{1,0}(X) = 1$.
- * Since $h^{1,0}(X) \neq 0$, X is **not rational**.

Example (Surfaces (dim 2))

A smooth cubic $X \subset \mathbb{P}^3$ is rational.

Intermediate Jacobian and Cubic Threefolds

Let $X \subset \mathbb{P}^4$ be a smooth cubic threefold.

Intermediate Jacobian (Clemens-Griffiths)

For X with $\dim_{\mathbb{C}} X = 3$:

$$J^2(X) := \frac{(H^{2,1})^*}{H^3(X, \mathbb{Z})}$$

Key properties:

- * Principally polarized abelian variety of dimension $\frac{1}{2} b_3(X) = 5$
- * Obstruction to rationality: $J^2(X)$ is **not** isomorphic to $\text{Jac}(C)$ for any curve C

Example (Hodge Diamond)

$$\begin{array}{ccc} & & 1 \\ & & 1 \\ & 5 & & 5 \\ & & 1 \\ & & 1 \end{array}$$

For cubic threefolds: $h^{p,q} = 0$ if $p \neq q$ and $p + q \neq 3$.

Theorem (Clemens-Griffiths (1972))

A smooth cubic threefold $X \subset \mathbb{P}^4$ is irrational because:

- * Its intermediate Jacobian $J^2(X)$ is the Jacobian of a curve **only if** X is a blowup of \mathbb{P}^3 .
- * For cubic threefolds, $J^2(X)$ decomposes as a product of Jacobians of curves $\iff X$ is rational.

Key Computation

For $X \subset \mathbb{P}^4$:

$$H^{2,1}(X) \cong H^{1,2}(X) \cong \mathbb{C}^5, \quad H^3(X, \mathbb{Z}) \cong \mathbb{Z}^{10}$$

Thus, $J^2(X)$ is a 5-dimensional abelian variety not isogenous to any product of curve Jacobians.

- * **Hodge-Theoretic Methods**
 - * Intermediate Jacobian (Clemens-Griffiths).
 - * Brauer group (Artin-Mumford).
- * **Geometric Methods**
 - * Birational automorphisms (Iskovskikh-Manin).
 - * Degenerations (Voisin, Kollár, Pirutka).
- * **Analytic Methods**
 - * Multiplier ideal sheaves (Nadel, Ein-Lazarsfeld).

Example: Cubic Fourfold (dim 4)

Let $X \subset \mathbb{P}^5$ be a smooth cubic fourfold.

- * Hodge diamond:

$$\begin{array}{ccccc} & & & & 1 \\ & & & & 1 \\ & & & & 1 \\ 1 & & 21 & & 1 \\ & & & & 1 \\ & & & & 1 \end{array}$$

- * Is X rational? Katzarkov-Kontsevich-Pantev-Yu: **no**.

Homological Mirror Symmetry

Statement: For a smooth projective variety X , HMS relates:

- * **B-model:** Derived category $D_{\text{coh}}^b(X)$.
- * **A-model:** Fukaya-Seidel category $FS(Y, W)$.

Example (Mirror of \mathbb{P}^2)

$$D^b(\mathbb{P}^2) \longleftrightarrow FS\left(\mathbb{C}^2, W = x + y + \frac{1}{xy}\right)$$

Hodge Structures in HMS

B-model (de Rham cohomology)


$$\text{for } \mathbb{P}^2 : \begin{array}{cccc} & & 1 & \\ & 0 & & 0 \\ & & 1 & 0 \\ 0 & & & 0 \\ & 0 & & 1 \end{array}$$

for the blow-up of \mathbb{P}^2 in a point : 0

$$\begin{array}{cccc} & & 1 & \\ 0 & & & 0 \\ & 2 & & 0 \\ 0 & & 0 & \\ & 1 & & \\ & & 1 & \\ 0 & & 0 & \\ \text{for the blow-up of } \mathbb{P}^2 \text{ in 6 points : } & 0 & 7 & 0 \\ & 0 & & 0 \\ & & 1 & \end{array}$$

for the blow-up of \mathbb{P}^2 in 6 points : 0

A-model (LG mirror)



ncHodge structure¹ on periodic cyclic homology links A/B-models.

¹Non-commutative Hodge structure encoding GW invariants.

Monodromy Operator and Rationality

Let $X \rightarrow Z_t$ be a family of smooth threefolds.

Monodromy Action

The monodromy operator acts on $H^2(Z_t, \mathbb{Z})$:

$$\mu : H^2(Z_t, \mathbb{Z}) \rightarrow H^2(Z_t, \mathbb{Z}), \quad \mu = \text{diag}(1, \epsilon, \epsilon^2), \quad \epsilon^3 = 1.$$

Example (Cubic Threefold)

For $X \subset \mathbb{P}^4$, μ is non-nilpotent $\implies X$ is irrational (Clemens-Griffiths).

Theorem (Katzarkov-Przyjalkowski)

Let X be a smooth Fano threefold with $\text{Pic}(X) \cong \mathbb{Z}$ and $X \not\cong \mathbb{P}^3$. Then X is rational if and only if the monodromy operator μ is nilpotent.

Three-Dimensional Cubic: LG Mirror

The LG mirror of a cubic threefold $X \subset \mathbb{P}^4$ is the fibration by open K3 surfaces given by the potential :

$$W = \frac{(x + y + z)^3}{xyz} + z \quad \text{on } \mathbb{C}^3.$$

This family of K3 surfaces has 3 singular fibers - two fibers with ordinary double points and one open-book singularity.



Let \mathcal{F} be the perverse sheaf of vanishing cycles of the potential. Then,

$$\dim \mathbb{H}^1(\mathcal{F}) = 5, \quad \dim \mathbb{H}^2(\mathcal{F}) = 4, \quad \dim \mathbb{H}^3(\mathcal{F}) = 5.$$

$A+B$

$H_{dR} + \text{Eingevalues of Quantum Multiplication}$

Splitting the Hodge Structure


If we further split the cohomology of X into generalized eigenspaces for the operator K of quantum multiplication by $c_1(X)$, or equivalently split the cohomology of \mathcal{F} according to the critical values of the potential, we obtain as a piece a Hodge structure \mathcal{H} which is exactly the Clemens-Griffiths invariant:



critical values of LG model

= eigenvalues of K : x_1, x_2, x_3

Splitting of $H^*(X) = \mathcal{H} + (1) + (1)$



Quantum differential equation:

$$\left(\frac{\partial}{\partial u} - \frac{1}{u^2} K + \frac{1}{u} G \right) \psi(u) = 0$$

- * K: Quantum multiplication by $c_1(X)$.
- * G: Connection matrix (flat coordinates).

Theorem (Katzarkov-Kontsevich-Pantev-Yu)

For a projective variety X :

- * **Decomposition:** $H^*(X)$ splits into H_{λ_i} , labeled by eigenvalues of K .
- * **Birational invariance:** Elementary pieces H_{λ_i} (modulo codimension ≥ 2) are birational invariants.

Applications

- * Singular fibers of LG mirror \leftrightarrow eigenvalues of K .
- * Integral Hodge structure on $\mathbb{H}^i(\mathcal{F})$ is computable via $\text{QH}(X)$.

Atoms and Euler Fields

Hodge Subspace and Euler Field

- * Let X be a complex projective variety. Consider the subspace of even Hodge classes:

$$H_{\text{Hodge}}(X) := \bigoplus_i \left(H^{i,i}(X) \cap H^{2i}(X, \mathbb{Q}) \right)$$

- * This defines a purely even Frobenius manifold \mathcal{F}_X over \mathbb{K} .
- * The *Euler field* $Eu \in \Gamma(\mathcal{F}_X, T_{\mathcal{F}_X})$ is:

$$Eu = c_1(T_X) + \sum_{\substack{i \\ \deg \Delta_i \neq 2}} \frac{\deg \Delta_i - 2}{2} t_i \Delta_i$$

- * At a generic $p \in \mathcal{F}_X$, the spectrum of $Eu \star \cdot$ gives a μ -fold spectral cover.

Definition (Atoms)

Atoms $_X$ are the connected components of this spectral cover.

- * *Key Example:* If K_X is nef, Atoms $_X$ is trivial (quantum product preserves $H^{\geq \bullet}(X)$).

Asymptotics of the Quantum Differential Equation

Quantum differential equation (QDE):

$$\left(\frac{\partial}{\partial u} - \frac{K}{u^2} + \frac{G}{u} \right) \psi = 0$$

Eigenvalues: Asymptotic solutions $\psi(u) \sim e^{\sigma/u}$ correspond to eigenvalues σ of $K \star \cdot$.

Theorem (Non-rationality criterion)

Let X be a Fano hypersurface of degree d in \mathbb{P}^{N-1} . Define:

$$\delta := \dim X - 2 \cdot \frac{N-d}{d}.$$

If $\delta > \dim X - 2$, then X is not rational.

Example (4D Quartic)

$X \subset \mathbb{P}^5$, $d = 4$, $N = 6$:

$$\delta = 4 - 2 \cdot \frac{6-4}{4} = 3 > 2 \quad (\dim X - 2 = 2) \implies \text{not rational.}$$


Example (5D quartic)

$$\delta = 5 - 2 \left(\frac{7-4}{4} \right) = 5 - 3 \frac{1}{2} > 3$$

Example (3D cubic)


For the three-dimensional generic cubic:

$$\delta = 3 - 2 \left(\frac{5-3}{3} \right) = \frac{5}{3} > 3 - 2 \implies \text{not rational.}$$





2-Dimensional Cubic X_k

X_k a 2-dimensional cubic with $\text{Pic}(X_k) \cong \mathbb{Z}_2$.



$$|2K_{\mathbb{P}^1} + 5| \neq 0$$

On the LG side: $H^0 + H^2 + H^4 = \mathbb{Z}^9$, Steenbrink=0.



Example: Hypersurface Model

Setup:

- * Let $X_{\text{geom}} \subset (\mathbb{P}^1)^4$ be a smooth hypersurface of degree $(1, 1, 1, 1)$ over an algebraically closed field k .
- * **Key Fact:** X_{geom} is the blowup of $(\mathbb{P}^1)^3$ at an elliptic curve E .

Atomic Structure

- * 8 simple "point-like" atoms.
- * 1 atom α_E linked to E .

New Setup:

- * Define X over a non-closed field k .
- * Assume the Galois group mixes the 4 factors of $(\mathbb{P}^1)^4$.

Key Calculation

At the "naive" point $q_i = 1, t_j = 0$:

$$\text{Eigenvalues of } Eu \star \cdot = \left\{ \underbrace{\lambda_1}_{\text{mult } 1}, \underbrace{\lambda_2}_{\text{mult } 4}, \underbrace{\lambda_3}_{\text{mult } 7} \right\}$$

- * The third piece has Hodge numbers: 5 (middle), 1 (top/bottom).
- * Only 2 algebraic cycles defined over k .

Thank you!

